

Anti-12 Lipoygenase/ALOX12 Antibody Picoband™

Catalog Number: A02275-1

About ALOX12

ALOX12, Arachidonate 12-lipoxygenase, is an enzyme that in humans is encoded by the ALOX12 gene. By fluorescence in situ hybridization, the ALOX12 gene is located in band 17p13.1. The gene consists of 14 exons with 13 introns and spans approximately 15 kb of DNA. Arachidonate 12-lipoxygenase introduces a molecular oxygen into the C-12 position of arachidonic acid to produce 12-(S)-hydroperoxy-5,8,10,14-eicosatetraenoic acid. The major pathway of arachidonic acid metabolism in human platelets proceeds via a 12-lipoxygenase enzyme. Expression of the ALOX12 gene was detected in human erythroleukemia cells, platelets, and human umbilical vein endothelial cells by reverse transcription-PCR analysis.

Overview

Product Name	Anti-12 Lipoygenase/ALOX12 Antibody Picoband™
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-12 Lipoygenase/ALOX12 Antibody Picoband™ catalog # A02275-1. Tested in WB applications. This antibody reacts with Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P18054

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human 12 Lipoygenase, different from the related mouse sequence by six amino acids, and from the related rat sequence by seven amino acids.
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Mouse, Rat</p>

Anti-12 Lipoxygenase/ALOX12 Antibody Picoband™ (A02275-1) Images




Figure 1. Western blot analysis of 12 Lipoxygenase using anti-12 Lipoxygenase antibody (A02275-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

lane 1: rat spleen tissue lysates,

lane 2: mouse spleen tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-12 Lipoxygenase antigen affinity purified polyclonal antibody (Catalog # A02275-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for 12 Lipoxygenase at approximately 80KD. The expected band size for 12 Lipoxygenase is at 75-80KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-12 Lipoxygenase/ALOX12 Antibody™