

Anti-IDUA Antibody Picoband® Fluoro488 Conjugated

Catalog Number: A02281-2-Fluoro488

About IDUA

This gene encodes an enzyme that hydrolyzes the terminal alpha-L-iduronic acid residues of two glycosaminoglycans, dermatan sulfate and heparan sulfate. This hydrolysis is required for the lysosomal degradation of these glycosaminoglycans. Mutations in this gene that result in enzymatic deficiency lead to the autosomal recessive disease mucopolysaccharidosis type I (MPS I).

Overview

| | |
|----------------------|--|
| Product Name | Anti-IDUA Antibody Picoband® Fluoro488 Conjugated |
| Reactive Species | Human, Mouse, Rat |
| Application | Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, IHC, WB). Customers may select suitable applications according to their experimental needs. |
| Clonality | Polyclonal |
| Formulation | Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ . |
| Storage Instructions | At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light. |
| Host | Rabbit |
| Uniprot ID | P35475 |

Technical Details

| | |
|---------------------|--|
| Immunogen | E.coli-derived human IDUA recombinant protein (Position: A28-D634). Human IDUA? shares?80.1%? amino acid?(aa)? sequence? identity? with mouse? IDUA. |
| Cross Reactivity | No cross reactivity with other proteins. |
| Isotype | IgG |
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Purification | Immunogen affinity purified. |
| Conjugate | Fluoro488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm |
| Suggested Dilutions | Optimal dilutions should be determined by end users. |

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IDUA Antibody - Fluoro488

For Research Use Only. Not for use in diagnostic procedures.