

Anti-Mannose Receptor/MRC1 Picoband® Antibody

Catalog Number: A02285-2

About MRC1

The mannose receptor (Cluster of Differentiation 206, CD206) is a C-type lectin primarily present on the surface of macrophages, immature dendritic cells and liver sinusoidal endothelial cells, but is also expressed on the surface of skin cells such as human dermal fibroblasts and keratinocytes. It is mapped to 10p12.33. The recognition of complex carbohydrate structures on glycoproteins is an important part of several biological processes, including cell-cell recognition, serum glycoprotein turnover, and neutralization of pathogens. The protein encoded by this gene is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. The protein has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment.

Overview


Product Name	Anti-Mannose Receptor/MRC1 Picoband® Antibody
Reactive Species	Human, Monkey, Mouse, Rat
Description	Boster Bio Anti-Mannose Receptor/MRC1 Picoband® Antibody catalog # A02285-2. Tested in ELISA, IF, IHC, WB applications. This antibody reacts with Human, Monkey, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IF, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P22897

Technical Details


Immunogen	E.coli-derived human Mannose Receptor/MRC1 recombinant protein (Position: D21-A1140).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti- Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human, Monkey, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human, Mouse, Rat Immunofluorescence, 5 ug/ml, Human, Mouse ELISA, 0.1-0.5ug/ml, -


Anti-Mannose Receptor/MRC1 Picoband® Antibody (A02285-2) Images



Western blot analysis of MRC1 using anti-MRC1 antibody (A02285-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat lung tissue lysates, Lane 2: mouse lung tissue lysates, Lane 3: human liver tissue lysates, Lane 4: monkey lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MRC1 antigen affinity purified polyclonal antibody (Catalog # A02285-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MRC1 at approximately 190-200 kDa. The expected band size for MRC1 is at 166 kDa.




IHC analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


IHC analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat




serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of rat liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IF analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. DyLight®550 Conjugated Goat Anti-Rabbit IgG (BA1135) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IF analysis of MRC1 using anti-MRC1 antibody (A02285-2). MRC1 was detected in a paraffin-embedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-MRC1 Antibody (A02285-2) overnight at 4°C. DyLight®550 Conjugated Goat Anti-Rabbit IgG (BA1135) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.


CS attenuates MMT in RIF rats. (A) Co-staining of the alpha-SMA (green) with CD68 (red) by Double Immunofluorescence. (B) Co-localization analysis of CD68+alpha-SMA + MMT cells. (C) Staining intensity of CD68+alpha-SMA + MMT cells was quantified. (D) Co-staining of the alpha-SMA (green) with CD206 (red) by




Double Immunofluorescence. (E) Co-localization analysis of CD206+alpha-SMA + MMT cells. (F) Staining intensity of CD206+alpha-SMA + MMT cells was quantified (n = 4). (G) mRNA Expression of AKT1, EGFR, IL-6, and IL-10 in RIF rats kidney (n = 3). ## p < 0.01 vs. the sham group; * p < 0.05 vs. the model group; ** p < 0.01 vs. the model group. Scale bars (40 um, 10 um).Index in PubMed under a CC BY license. PMID: 41142245



Wound healing with different hydrogels in vivo . (A) Preparation and observation of diabetic chronic wounds related to orthopedics. (B) Photographs of the wound treated with the hydrogel in different hydrogel groups. (C) Wound recovery curve of different hydrogel groups. (D) H&E staining and Masson staining of wound tissue on day 10 in different hydrogel groups. (E) IHC staining of wound tissue in iNOS and CD206.Index in PubMed under a CC BY license. PMID: 41190287




Hydrogel-induced macrophage M2 polarization in vitro . (A) Flow cytometry analysis of the macrophage surface marker CD206. (B) Immunofluorescence of CD206 in macrophages treated with different hydrogels. (C) Quantitative analysis of flow cytometry in CD206. (D) Mean fluorescence intensity of CD206 in macrophages treated with different hydrogels. (E, F) Quantitative analysis and immunofluorescence of CD163 in macrophages treated with different hydrogels.Index in PubMed under a CC BY license. PMID: 41190287




IL-1beta-primed ERCs reduced macrophage infiltration and inflammatory cytokine expressions in colons. Intra-colon macrophage infiltration was evaluated by immunohistochemical staining. Specifically, we stained iNOS for detecting M1 cell infiltration, and CD206 for measuring M2 cell infiltration. The represent IHC images of mouse colons and quantitative data for cell counts of each group are shown in A (x 200) and B , respectively. The concentrations of inflammatory cytokine productions in colonic tissues were determined by ELISA kit and the relative mRNA expression changes were performed by real-time PCR. IFN-gamma (C , D), IL-17 (E , F), TNF-alpha (G , H), IL-6 (I , J), IL-4 (K , L), and IL-10 (M , N) were shown, respectively, which are majorly secreted by Th1, Th2, Th17 cells, or CD206 + macrophages, and closely associated with the development of UC. Data were presented as mean ± standard deviation, and p values were calculated by using one-way ANOVA followed by the least significant difference (LSD) test (n = 6, * p < 0.05, ** p < 0.01, *** p < 0.001) Index in PubMed under a CC BY license. PMID: 34090510


IL-1beta-primed ERCs reduced the population of mature




DCs, but increased M2 macrophages in colitis mice. To determine whether each treatment has an influence on regulating DC and macrophage phenotypes, anti-CD11c antibody and antigen presenting-related antibodies (anti-MHCII, anti-CD86) were used to measure mature DCs, while anti-CD68 antibody and anti-CD206 antibody were used for M2 phenotype macrophages in spleens. A Representative dot plots of CD11c + MHCII + DCs and CD11c + CD86 + DCs in spleens. B Dot plots of CD68 + CD206 + macrophages. C - E Percentage of CD11c + MHCII + DCs, CD11c + CD86 + DCs, and CD68 + CD206 + macrophages, respectively. Data were mean \pm SD (n = 6, * p < 0.05, ** p < 0.01, *** p < 0.001). p values were analyzed by one-way ANOVA followed by the LSD test Index in PubMed under a CC BY license. PMID: 34090510



Identification of the isolated polarized M2 microglia and exosomes. (A) The expression of M2 microglia markers Arg1 and CD206 detected by RT-PCR. (B) The expression of M2 microglia markers Arg1 and CD206 detected by Western blot analysis. (C) The expression of M2 microglia markers Arg1 and CD206 detected by immunofluorescence. (D) Images of M2-EXOs detected using TEM. (E) Measurements of particle sizes range of the isolated exosomes using NTA. (F) Detection of exosomal marker proteins CD63 and TSG101 using western blot. All experiments were repeated three times. *** P < 0.001 vs. control. Index in PubMed under a CC BY license. PMID: 35573832




Histological staining evaluation. FU (FA-UIO-66-NH₂), BFU (Bai@FA-UIO-66-NH₂). (A) H&E staining and (B) Safranin O staining of the knee joint of SD rats and (C) partial magnified detail (bar = 200 μ m). (D) Immunofluorescence staining of iNOS/CD206/DAPI of the synovial of SD rats (scale bar = 200 μ m) Index in PubMed under a CC BY license. PMID: 38724958




In vitro anti-inflammatory effect produced by driving the polarization of macrophage M1 towards M2. FU (FA-UIO-66-NH₂), BFU (Bai@FA-UIO-66-NH₂). (A-G) In vitro mRNA expression of IL-1beta, IL-6, iNOS, Arg-1, CD86, IL-10 and CD206. (H-J) Immunofluorescence staining of iNOS and CD206 after different treatments. (H) (scale bar = 100 μ m) and Corresponding fluorescence intensity quantification (I-J). Flow cytometry analysis of specific marker of M1 macrophage iNOS (K) and M2 macrophage CD206 (L). (n = 3, mean \pm SD, “*” symbol compared with normal group, * P


Mice reconstituted with FGF2 KO macrophages and subjected to CLP demonstrate increased M1 polarization in lung tissue. a - f The presence and levels of CD206, CD86, and F4/80 markers on macrophages within lung tissue were identified and quantitatively assessed using




immunofluorescence staining. Bar is 20 μ m. * p



Effect of FGF2 deficiency on BMDM apoptosis and polarization. a - c FGF2 deletion increased BMDM apoptosis. a Apoptosis in BMDM deprived of FBS for 24 h was assessed by flow cytometry (n = 4). b - c Percentage of PI + Annexin V + and PI- Annexin V + BMDM after starvation. d - k FGF2 deletion in BMDM promoted M1 polarization. d - g Flow cytometric analysis of macrophage markers in BMDM treated with LPS or IL4, including CD86, iNOS, CD206, and Arg1 (n = 3). h - k The levels of CD86, iNOS, CD206 and Arg1 in BMDM after treatment with LPS or IL4. N represents no treatment; * p



YNJ reduces the secretion of inflammatory factors and promotes M2 macrophage polarization. (A-C) ELISA of TNF-alpha (A) , IL-6 (B) , and IL-1beta (C) contents in the supernatant of cultured RAW264.7 macrophages. (D) IF images of CD86 and CD206 detection in cultured RAW264.7 macrophages. (E, F) Statistical analysis of CD86 and CD206 expression by IF. The data are presented as the mean \pm SD (n = 6). ** P < 0.01. Index in PubMed under a CC BY license. PMID: 39925847



PF-127/hADSCs-Exos complex treatment inhibits inflammatory reaction. a Representative images of TNF-alpha immunostaining at 4, 7, and 10 days after treatment. Scale bar = 20 μ m. b Quantification of TNF-alpha + IHC stained tissues. c Representative images illustrating IHC results of IL-6 at 4, 7, and 10 days after surgery. Scale bar = 20 μ m. d Quantification of IL-6 + IHC stained tissues. e IHC images of wound sections stained with CD68 on days 4, 7, and 10 post-wounding. Scale bar = 20 μ m. f Quantification of the number of CD68 positive cells in the wound area on days 4, 7, and 10. g IHC images of wound sections stained with CD206 at days 4, 7, and 10 post-wounding. Scale bar = 20 μ m. h Quantification of the number of CD206 positive cells in the wound area on days 4, 7, and 10. In b, d, and f , data are shown as mean \pm SEM; n = 6 for each group. * p

8 Publications Citing This Product

1. PubMed ID: 10.1186/s13287-021-02392-9, IL-1beta pre-stimulation enhances the therapeutic effects of endometrial regenerative cells on experimental colitis

2. PubMed ID: 10.1155/2020/2462039, Protective Effects of Two Safflower Derived Compounds, Kaempferol and Hydroxysafflor Yellow A, on Hyperglycaemic Stress-Induced Podocyte Apoptosis via Modulating of Macrophage M1/M2 Polarization

3. PubMed ID: 10.1155/2020/8263598, Protective Effects of Thalidomide on High-Glucose-Induced Podocyte Injury through In Vitro Modulation of Macrophage M1/M2 Differentiation

Visit bosterbio.com/anti-mannose-receptor-mrc1-picoband-trade-antibody-a02285-2-boster.html to see all 8 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Mannose Receptor/MRC1 Antibody

For Research Use Only. Not for use in diagnostic procedures.