

Anti-CD204/Msr1 Antibody Picoband®

Catalog Number: A02349-2

About Msr1

Macrophage scavenger receptor 1, also known as MSR1, is a protein which in humans is encoded by the MSR1 gene. Enables amyloid-beta binding activity; cargo receptor activity; and low-density lipoprotein particle binding activity. Involved in several processes, including endocytosis; positive regulation of cholesterol storage; and positive regulation of macrophage derived foam cell differentiation. Acts upstream of or within lipoprotein transport. Located in plasma membrane. Is expressed in several structures, including alimentary system; limb; neural tube; reproductive system; and yolk sac. Human ortholog(s) of this gene implicated in Barrett's esophagus; arteriosclerosis; and prostate cancer. Orthologous to human MSR1 (macrophage scavenger receptor 1).

Overview


Product Name	Anti-CD204/Msr1 Antibody Picoband®
Reactive Species	Mouse
Description	Boster Bio Anti-CD204/Msr1 Antibody Picoband® catalog # A02349-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P30204

Technical Details


Immunogen	E.coli-derived mouse CD204/Msr1 recombinant protein (Position: M1-D451).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Mouse ELISA, 0.1-0.5 ug/ml, -

Anti-CD204/Msr1 Antibody Picoband® (A02349-2) Images



Western blot analysis of CD204/Msr1 using anti-CD204/Msr1 antibody (A02349-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: mouse RAW264.7 whole cell lysates, Lane 2: mouse ANA-1 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD204/Msr1 antigen affinity purified polyclonal antibody (Catalog # A02349-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CD204/Msr1 at approximately 75-90 kDa. The expected band size for CD204/Msr1 is at 71 kDa.



Flow Cytometry analysis of RAW264.7 cells using anti-CD204/Msr1 antibody (A02349-2). Overlay histogram showing RAW264.7 cells stained with A02349-2 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-CD204/Msr1 Antibody (A02349-2, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD204/Msr1 Antibody

For Research Use Only. Not for use in diagnostic procedures.