

Anti-FABP2/I-FABP Antibody Picoband®

Catalog Number: A02378

About Fabp2

FABP 2, Fatty acid-binding protein 2, is a protein that in humans is encoded by the FABP2 gene. Using a human cDNA probe, the gene is assigned to chromosome 4 in somatic cell hybrids. FABP 2 gene contains four exons and is an abundant cytosolic protein in small intestine epithelial cells. The FABPs belong to a multigene family with nearly twenty identified members. And FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. Also, they may be responsible in the modulation of cell growth and proliferation.

Overview


Product Name	Anti-FABP2/I-FABP Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-FABP2/I-FABP Antibody Picoband® catalog # A02378. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P55050

Technical Details


Immunogen	E.coli-derived mouse FABP2/I-FABP recombinant protein (Position: A2-E132). Mouse FABP2/I-FABP shares 77.9% and 92.4% amino acid (aa) sequence identity with human and rat FABP2/I-FABP, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Mouse, Rat Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat


Anti-FABP2/I-FABP Antibody Picoband® (A02378) Images



Western blot analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat liver tissue lysates, Lane 2: rat small intestine tissue lysates, Lane 3: mouse liver tissue lysates, Lane 4: mouse small intestine tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FABP2/I-FABP antigen affinity purified polyclonal antibody (Catalog # A02378) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FABP2/I-FABP at approximately 15 kDa. The expected band size for FABP2/I-FABP is at 15 kDa.




IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378). FABP2/I-FABP was detected in paraffin-embedded section of mouse intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378). FABP2/I-FABP was detected in paraffin-embedded section of rat intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378). FABP2/I-FABP was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer



(pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-FABP2/I-FABP Antibody

For Research Use Only. Not for use in diagnostic procedures.