

# **Anti-Bag-1 Antibody**

Catalog Number: A02423-1

#### Overview


| Product Name         | Anti-Bag-1 Antibody  |
|----------------------|--|
| Reactive Species     | Canine, Human, Mouse, Rat  |
| Description          | Boster Bio Anti-Bag-1 Antibody (Catalog# A02423-1). Tested in IP, IHC, WB application(s). This antibody reacts with Canine, Rat, Mouse, Human. |
| Application          | IP, IHC, WB  |
| Clonality            | Polyclonal   |
| Formulation          | 50 ul sera   |
| Storage Instructions | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.                  |
| Host                 | Rabbit   |
| Uniprot ID           | Q99933   |

## **Technical Details**


| Immunogen                  | A synthetic peptide of Bag-1 protein (amino acids 205-219 QETERLQSTNLALAE)  |
|----------------------------|---|
| Predicted Reactive Species | Mouse, Rat  |
| Isotype                    | Rabbit IgG  |
| Form                       | Liquid  |
| Concentration              | 0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.  |
| Suggested Dilutions        | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  WB: 1:1000-1:2000  IHC (paraffin): 1:1000-1:5000  IHC (frozen): Users should optimize  IP: 1:50-1:200  IF/ICC: 1:500-1:2000 |



## Anti-Bag-1 Antibody (A02423-1) Images



Formalin-fixed, paraffin-embedded section of human colon stained for Bag-1 using primary antibody dilution ratio of 1:2000. Hematoxylin-eosin counterstain.



Formalin-fixed, paraffin-embedded sections of human colorectal cancer stained for Bag-1 using primary antibody dilution ratio of 1:2000. Hematoxylin-eosin counterstain. Samples from two patients (A and B) are shown. A1 and B1 are higher magnifications from A and B, respectively.

#### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Bag-1 Antibody