

## Anti-C19orf80/ANGPTL8 Antibody Picoband™

Catalog Number: A02471

### About ANGPTL8

ANGPTL8 (also known as lipasin, originally Betatrophin) is a protein that in humans is encoded by the C19orf80 gene. The gene for betatrophin lies on mouse chromosome 9 (gene symbol: Gm6484) and on human chromosome 19 (gene symbol: C19orf80). The encoded 22 kDa protein contains an N-terminal secretion signal and two coiled-coil domains and is a member of the angiopoietin-like (ANGPTL) protein family. However, in contrast to other ANGPTL proteins, betatrophin lacks the C-terminal fibrinogen-like domain, and therefore it is an atypical member of the ANGPTL family. It shares with Angptl4 and Angptl3 the ability to inhibit the enzyme Lipoprotein lipase (LPL), and its hepatic overexpression causes elevation of circulating Triglyceride levels in mice. In mice betatrophin is secreted by the liver. It was hoped that betatrophin or its homolog in humans may provide an effective treatment for type 2 diabetes and perhaps even type I diabetes.

### Overview


Product Name	Anti-C19orf80/ANGPTL8 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-C19orf80/ANGPTL8 Antibody Picoband™ catalog # A02471. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q6UXH0

### Technical Details

Immunogen	E.coli-derived human C19orf80 recombinant protein (Position: A22-A198). Human C19orf80 shares 74% amino acid (aa) sequence identity with mouse C19orf80.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>

## Anti-C19orf80/ANGPTL8 Antibody Picoband™ (A02471) Images



**Figure 1. Western blot analysis of C19orf80 using anti-C19orf80 antibody (A02471).**

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HepG2 whole Cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-C19orf80 antigen affinity purified polyclonal antibody (Catalog # A02471) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for C19orf80 at approximately 22KD. The expected band size for C19orf80 is at 22KD.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-C19orf80/ANGPTL8 Antibody™