

Anti-liver FABP/Fabp1 Antibody Picoband®

Catalog Number: A02477

About Fabp1

Fatty acid binding protein 1, liver, also known as FABP1 or FABPL, is a human gene locating at 2p11. FABP1 encodes the fatty acid binding protein found in liver. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind free fatty acids, their CoA derivatives, bilirubin, organic anions, and other small molecules. FABP1 and FABP6 (the ileal fatty acid binding protein) are also able to bind bile acids. It is thought that FABPs roles include fatty acid uptake, transport, and metabolism. The liver form of FABP may be identical to the major liver protein-1 (Lvp-1), which is encoded by a gene situated within 1 cM of Ly-2.

Overview


Product Name	Anti-liver FABP/Fabp1 Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-liver FABP/Fabp1 Antibody Picoband® catalog # A02477. Tested in ELISA, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P02692

Technical Details

Immunogen	E. coli-derived rat liver FABP recombinant protein (Position: M1-I127).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot,0.1-0.5ug/ml ELISA,0.1-0.5ug/ml

Anti-liver FABP/Fabp1 Antibody Picoband® (A02477) Images



Western blot analysis of Liver FABP/Fabp1 using anti-Liver FABP/Fabp1 antibody (A02477). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat liver tissue lysates, Lane 2: rat liver tissue lysates, Lane 3: mouse liver tissue lysates, Lane 4: mouse liver tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Liver FABP/Fabp1 antigen affinity purified polyclonal antibody (Catalog # A02477) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Liver FABP/Fabp1 at approximately 14 kDa. The expected band size for Liver FABP/Fabp1 is at 14 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-liver FABP/Fabp1 Antibody

For Research Use Only. Not for use in diagnostic procedures.