

## Anti-HSD17B2 Antibody Picoband® Fluoro488 Conjugated

Catalog Number: A02506-1-Fluoro488

### About HSD17B2

HSD17B2(17-BETA-HYDROXYSTEROID DEHYDROGENASE II), also called 17-BETA-HSD II, is an enzyme which has 387 amino acids with a predicted molecular weight of 42,782 and associates with the membranes of the endoplasmic reticulum. Its cytogenetic location is 16q23.3. The type 2 enzyme was capable of catalyzing the interconversion of testosterone and androstenedione, as well as estradiol and estrone. HSD17B2 mRNA was detected in 18 of 42(43%) adenomas but not in prolactinomas. In the human endometrium, inactivation of 17-beta-estradiol to estrone is catalyzed by HSD17B2. And HSD17B2 activity distinguishes between disease-free and diseased endometria. HSD17B2 efficiently catalyzes the oxidative metabolism of androgens and estrogens, and it is expressed in a large series of human peripheral tissues. The previous paradigm that HSD17B2 activity in the endometrium is elevated during the secretory phase is confined to diseased endometrium but not to disease-free endometrium.

### Overview

Product Name	Anti-HSD17B2 Antibody Picoband® Fluoro488 Conjugated
Reactive Species	Human, Mouse
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% NaN <sub>3</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P37059

### Technical Details

Immunogen	E.coli-derived human HSD17B2 recombinant protein (Position: Q75-R365). Human HSD17B2 shares 63.6% and 65.2% amino acid (aa) sequence identity with mouse and rat HSD17B2, respectively.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	Fluoro488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HSD17B2 Antibody - Fluoro488

For Research Use Only. Not for use in diagnostic procedures.