BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Livin/Birc7 Antibody Picoband®

Catalog Number: A02577-2

About Birc7

Baculoviral IAP repeat-containing protein 7 is a protein that in humans is encoded by the BIRC7 gene. This protein is a member of the family of inhibitor of apoptosis proteins(IAP) and contains a single copy of a baculovirus IAP repeat(BIR) and a RING finger motif. Its gene is mapped to 20q13.3. It has got 280- and 298- amino acids. The protein is highly expressed in melanomas while lowly expressed in some lymphomas, fetal kidney, fetal liver, testis and thymus. BIRC7 plays a vital role in blocking apoptosis induced.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Livin/Birc7 Antibody Picoband®
Reactive Species	Mouse
Description	Boster Bio Anti-Livin/Birc7 Antibody Picoband® catalog # A02577-2. Tested in ELISA, WB applications. This antibody reacts with Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	A2AWP0

Technical Details


Immunogen	E.coli-derived mouse Livin/Birc7 recombinant protein (Position: Q41-S285).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse ELISA, 0.1-0.5 ug/ml, -

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Livin/Birc7 Antibody Picoband® (A02577-2) Images



Western blot analysis of Livin/Birc7 using anti-Livin/Birc7 antibody (A02577-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: mouse heart tissue lysates, Lane 2: mouse lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Livin/Birc7 antigen affinity purified polyclonal antibody (Catalog # A02577-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Livin/Birc7 at approximately 33 kDa. The expected band size for Livin/Birc7 is at 33 kDa.

Anti-Livin/Birc7 Antibody

For Research Use Only. Not for use in diagnostic procedures.