

Anti-Six3 Antibody Picoband™

Catalog Number: A02604

About SIX3

Homeobox protein SIX3 is a protein that in humans is encoded by the SIX3 gene. This gene encodes a member of the sine oculishomeobox transcription factor family. The encoded protein plays a role in eye development. Mutations in SIX3 are the cause of a severe brain malformation, called holoprosencephaly type 2 (HPE2). In HPE2, the brain fails to separate into two hemispheres during early embryonic development, leading to eye and brain malformations, which result in serious facial abnormalities. A mutant zebrafish knockout model has been developed, in which the anterior part of the head was missing due to the atypical increase of Wnt1 activity. When injected with SIX3, these zebrafish embryos were able to successfully develop a normal forebrain. When SIX3 was turned off in mice, resulting in a lack of retina formation due to excessive expression of Wnt8b in the region where the forebrain normally develops. Both of these studies demonstrate the importance of SIX3 activity in brain and eye development.

Overview

Product Name	Anti-Six3 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Six3 Antibody Picoband™ catalog # A02604. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O95343

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Six3, different from the related mouse sequence by two amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human</p>

Anti-Six3 Antibody Picoband™ (A02604) Images




Figure 1. Western blot analysis of Six3 using anti-Six3 antibody (A02604).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat brain tissue lysate,

Lane 2: mouse brain tissue lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Six3 antigen affinity purified polyclonal antibody (Catalog # A02604) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Six3 at approximately 35KD. The expected band size for Six3 is at 35KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Six3 Antibody™