

Anti-53BP2/TP53BP2 Antibody Picoband®

Catalog Number: A02609-3

About TP53BP2

tumor protein p53 binding protein 2


Overview

Product Name	Anti-53BP2/TP53BP2 Antibody Picoband®
Reactive Species	Human, Mouse
Description	Boster Bio Anti-53BP2/TP53BP2 Antibody Picoband® catalog # A02609-3. Tested in WB, IP, ELISA applications. This antibody reacts with Human, Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IP, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q13625


Technical Details

Immunogen	E.coli-derived human 53BP2/TP53BP2 recombinant protein (Position: Q97-E1096).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse Immunoprecipitation, 0.5-2 ug/ml, Human ELISA, 0.1-0.5 ug/ml

Anti-53BP2/TP53BP2 Antibody Picoband® (A02609-3) Images



Western blot analysis of 53BP2/TP53BP2 using anti-53BP2/TP53BP2 antibody (A02609-3). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human THP-1 whole cell lysates, Lane 2: human SH-SY5Y whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human HEL whole cell lysates, Lane 5: mouse lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-53BP2/TP53BP2 antigen affinity purified polyclonal antibody (A02609-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for 53BP2/TP53BP2 at approximately 160 kDa. The expected band size for 53BP2/TP53BP2 is at 126 kDa.



Immunoprecipitating 53BP2/TP53BP2 in K562 whole cell lysate. Western blot analysis of 53BP2/TP53BP2 using anti-53BP2/TP53BP2 antibody (A02609-3). Lane 1: K562 whole cell lysates (30ug), Lane 2: Rabbit control IgG instead of anti-53BP2/TP53BP2 antibody in K562 whole cell lysate, Lane 3: anti-53BP2/TP53BP2 antibody (2ug) + K562 whole cell lysate (500ug). After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-53BP2/TP53BP2 antigen affinity purified polyclonal antibody (A02609-3) at a dilution of 0.5 ug/mL and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for 53BP2/TP53BP2 at approximately 160 kDa. The expected band size for 53BP2/TP53BP2 is at 126 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-53BP2/TP53BP2 Antibody

For Research Use Only. Not for use in diagnostic procedures.