

## Anti-JCHAIN Antibody Picoband® APC Conjugated

Catalog Number: A02644-2-APC

### About JCHAIN

Immunoglobulin J chain, also known as IGJ and IGCJ, is a secreted polypeptide which is the first immunoglobulin-related polypeptide expressed during the embryogenesis and differentiation of B cells in the fetal liver. The joining Immunoglobulin J chain is a small polypeptide, expressed by mucosal and glandular plasma cells, which regulates polymer formation of immunoglobulin (Ig)A and IgM. Immunoglobulin J chain / IGJ serves to link two monomer units of either IgM or IgA. In the case of IgM, the J chain-joined dimer is a nucleating unit for the IgM pentamer, and in the case of IgA, it induces larger polymers. Immunoglobulin J chain / IGJ also helps to bind these immunoglobulins to the secretory component. J-chain incorporation into polymeric IgA (pIgA, mainly dimers) and pentameric IgM endows these antibodies with several salient features. Immunoglobulin J chain / IGJ is involved in creating the binding site for pIgR / SC in the Ig polymers, not only by determining the polymeric quaternary structure but also by interacting directly with the receptor protein. Both the immunoglobulin J chain / IGJ and the pIgR/SC are key proteins in secretory immunity.

### Overview

Product Name	Anti-JCHAIN Antibody Picoband® APC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, IHC, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% Na <sub>3</sub> N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P01591

### Technical Details

Immunogen	E.coli-derived human JCHAIN recombinant protein (Position: Q23-D159). Human JCHAIN shares 77.4% and 73.5% amino acid (aa) sequence identity with mouse and rat JCHAIN, respectively.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-JCHAIN Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.