BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-CIDE-A Antibody

Catalog Number: A02671-1

About CIDEA

Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. DFF45/ICAD has been identified as inhibitor of caspase activated DNase DFF40/CAD. DFF45 related proteins CIDE-A and CIDE-B (for cell death-inducing DFF-like effector A and B) were recently identified. CIDE contains a new type of domain termed CIDE-N, which has high homology with the regulatory domains of DFF45/ICAD and DFF40/CAD. Expression of CIDE-A induces DNA fragmentation and activates apoptosis, which is inhibited by DFF45. CIDE-A is a DFF45-inhibitable effector that promotes cell death and DNA fragmentation. CIDE-A is expressed in many tissues.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-CIDE-A Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-CIDE-A Antibody (Catalog # A02671-1). Tested in ELISA, WB, IHC-P, IF applications. This antibody reacts with Human, Mouse.
Application	ELISA, IF, IHC-P, WB
Clonality	Polyclonal
Formulation	CIDE-A Antibody is supplied in PBS containing ≤0.09% sodium azide.
Storage Instructions	CIDE-A antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.
Host	Rabbit
Uniprot ID	O60543

Technical Details


Immunogen	CIDE-A antibody was raised against an 18 amino acid peptide near the carboxy terminus of mouse CIDE-A. The immunogen is located within the last 50 amino acids of CIDE-A.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Predicted Reactive Species	Bovine, Chicken, Mouse, Rat
Cross Reactivity	It has no cross activity to CIDE-B.
Isotype	IgG
Form	Liquid
Concentration	1 mg/mL
Purification	CIDE-A Antibody is affinity chromatography purified via peptide column.
Suggested Dilutions	CIDE-A antibody can be used for detection of CIDE-A by Western blot at 0.5 ug/mL. Antibody can also be used for immunohistochemistry starting at 2.5 ug/mL. For immunofluorescence start at 20 ug/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human and mouse samples and Immunofluorescence in human and mouse samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-CIDE-A Antibody (A02671-1) Images




Western blot analysis of CIDE-A in mouse heart tissue lysate with CIDE-A antibody at 1:500 dilution.



Immunohistochemistry of CIDE-A in mouse heart tissue with CIDE-A antibody at 5 ug/mL.




Immunofluorescence of CIDE-A in Mouse Heart cells with CIDE-A antibody at 20 ug/mL.




Immunohistochemistry of CIDE-A in human brain tissue with CIDE-A antibody at 2.5 ug/ml.

Immunofluorescence of CIDE-A in human brain tissue with CIDE A antibody at 20 ug/mL.Green: CIDE-A Antibody (A02671-1) Blue: DAPI staining

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com





Immunohistochemistry of CIDE-A in human brain tissue with CIDE A antibody at 2.5 ug/mL.

Anti-CIDE-A Antibody

For Research Use Only. Not for use in diagnostic procedures.