

Anti-BAFF Receptor/Tnfrsf13c Antibody Picoband™

Catalog Number: A02865-2

About Tnfrsf13c

Tumor necrosis factor receptor superfamily member 13C (TNFRSF13C), also known as BAFFR, is a protein in humans is encoded by the TNFRSF13C gene. The BAFFR gene is mapped to chromosome 22q13.1-q13.31. It has got 184 amino acid transmembrane protein which is 56% identical to the mouse protein. B cell-activating factor (BAFF) enhances B-cell survival in vitro and is a regulator of the peripheral B-cell population. BAFF plays a crucial role in B cell development and can function through receptors other than BCMA.

Overview

| | |
|----------------------|---|
| Product Name | Anti-BAFF Receptor/Tnfrsf13c Antibody Picoband™ |
| Reactive Species | Mouse, Rat |
| Description | Boster Bio Anti-BAFF Receptor/Tnfrsf13c Antibody Picoband™ catalog # A02865-2. Tested in ELISA, WB applications. This antibody reacts with Mouse, Rat. |
| Application | ELISA, WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q9D8D0 |

Technical Details

| | |
|-------------------------------|--|
| Immunogen | E. coli-derived mouse BAFF Receptor recombinant protein (Position: M1-A71). |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the |

optimal dilution ratio for your samples.
Some PubMed article(s) citing the expression level of this target are as follows:
Boster Bio's internal QC testing used:
Western blot, 0.1-0.5ug/ml
Direct ELISA, 0.1-0.5ug/ml

Anti-BAFF Receptor/Tnfrsf13c Antibody Picoband™ (A02865-2) Images




Figure 1. Western blot analysis of BAFF Receptor using anti-BAFF Receptor antibody (A02865-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat spleen tissue lysates,
Lane 2: rat thymus tissue lysates,
Lane 3: mouse spleen tissue lysates,
Lane 4: mouse thymus tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BAFF Receptor antigen affinity purified polyclonal antibody (Catalog # A02865-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for BAFF Receptor at approximately 19, 40KD. The expected band size for BAFF Receptor is at 19KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-BAFF Receptor/Tnfrsf13c Antibody™