

## Anti-ADAMTS4/ADAMTS4 Antibody Picoband®

Catalog Number: A02899

### About ADAMTS4

ADAMTS4, A disintegrin and metalloproteinase with thrombospondin motifs 4, is an enzyme that in humans is encoded by the ADAMTS4 gene. ADAMTS4 is a member of the large ADAMTS family of zinc-dependent proteases. The human ADAMTS4 gene is mapped to chromosome 1 by somatic cell hybrid analysis. The enzyme encoded by this gene lacks a C-terminal TS motif. It is responsible for the degradation of aggrecan, a major proteoglycan of cartilage, and brevican, a brain-specific extracellular matrix protein. The cleavage of aggrecan and brevican suggests key roles of this enzyme in arthritic disease and in the central nervous system, potentially, in the progression of glioma.

### Overview


Product Name	Anti-ADAMTS4/ADAMTS4 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-ADAMTS4/ADAMTS4 Antibody Picoband® catalog # A02899. Tested in ELISA, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O75173

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human ADAMTS4, different from the related mouse sequence by five amino acids, and from the related rat sequence by six amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human ELISA, 0.1-0.5ug/ml, -

## Anti-ADAMTS4/ADAMTS4 Antibody Picoband® (A02899) Images



Western blot analysis of ADAMTS4 using anti-ADAMTS4 antibody (A02899). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Recombinant Human ADAMTS4 Protein 0.5ng After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ADAMTS4 antigen affinity purified polyclonal antibody (Catalog # A02899) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ADAMTS4 at approximately 58KD. The expected band size for ADAMTS4 is at 53KD.

### 1 Publications Citing This Product

1. PubMed ID: 23990941, Electrospun Poly(L-lactide)/Poly(?-caprolactone) Blend Nanofibrous Scaffold: Characterization and Biocompatibility with Human Adipose-Derived Stem Cells

Visit [bosterbio.com/anti-adamts4-picoband-trade-antibody-a02899-boster.html](http://bosterbio.com/anti-adamts4-picoband-trade-antibody-a02899-boster.html) to see all 1 publications.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ADAMTS4/ADAMTS4 Antibody

For Research Use Only. Not for use in diagnostic procedures.