

Anti-CD105/ENG Antibody Picoband®

Catalog Number: A02997-1

About ENG

Endoglin (Osler-Rendu-Weber syndrome 1), CD105, is a type I membrane glycoprotein located on cell surfaces and is a part of the TGF beta receptor complex. Its gene is mapped to human chromosome 8. The protein consists of a homodimer of 180 kDA with disulfide links. It has been found on endothelial cells, activated macrophages, fibroblasts and smooth muscle cells. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling and has been found to be elevated in pregnant women who subsequently develop preeclampsia.

Overview


Product Name	Anti-CD105/ENG Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-CD105/ENG Antibody Picoband® catalog # A02997-1. Tested in Flow Cytometry, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P17813

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human CD105, different from the related mouse sequence by twelve amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human

Anti-CD105/ENG Antibody Picoband® (A02997-1) Images



Western blot analysis of CD105/ENG using anti-CD105/ENG antibody (A02997-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human SiHa whole cell lysates, Lane 2: human Hela whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD105/ENG antigen affinity purified polyclonal antibody (Catalog # A02997-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CD105/ENG at approximately 95 kDa. The expected band size for CD105/ENG is at 71 kDa.



Flow Cytometry analysis of U87 cells using anti-CD105/ENG antibody (A02997-1). Overlay histogram showing U87 cells stained with A02997-1 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-CD105/ENG Antibody (A02997-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

10 Publications Citing This Product

1. PubMed ID: -, Wu,H.,Jiang,X.,Li,Y.,Zhou,Y.,Zhang,T.,Zhi,P.,Gao,J.,Engineering Stem Cell Derived Biomimetic Vesicles for Versatility and Effective Targeted Delivery.Adv. Funct.Mater.2020, 30, 2006169.<https://doi.org/10.1002/adfm.202006169>
2. PubMed ID: 4002829, Antitumor activity of endogenous mFlt4 displayed on a T4 phage nanoparticle surface
3. PubMed ID: 28147322, VEGF promotes cartilage angiogenesis by phospho-ERK1/2 activation of Dll4 signaling in temporomandibular joint osteoarthritis caused by chronic sleep %u2026

Visit [bosterbio.com/anti-cd105-picoband-trade-antibody-a02997-1-boster.html](https://www.bosterbio.com/anti-cd105-picoband-trade-antibody-a02997-1-boster.html) to see all 10 publications.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD105/ENG Antibody

For Research Use Only. Not for use in diagnostic procedures.