

## Anti-Aquaporin 5/AQP5 Antibody Picoband®

Catalog Number: A03085

### About AQP5

Aquaporin 5, also known as AQP5, is a water channel protein. The aquaporins (AQPs) are a family of more than 10 homologous water transporting proteins expressed in many mammalian epithelia and endothelia. At least five AQPs are expressed in the eye: AQP0 (MIP) in lens fiber, AQP1 in cornea endothelium, ciliary and lens epithelia and trabecular meshwork, AQP3 in conjunctiva, AQP4 in ciliary epithelium and retinal Müller cells, and AQP5 in corneal and lacrimal gland epithelia. Among the seven human aquaporins cloned to date (AQPs 0-6), genes encoding the four most closely related aquaporins (AQP0, AQP2, AQP5, and AQP6) have been mapped to chromosome band 12q13, suggesting an aquaporin family gene cluster at this locus. Aquaporin 5 plays a role in the generation of saliva, tears and pulmonary secretions.

### Overview


Product Name	Anti-Aquaporin 5/AQP5 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Aquaporin 5/AQP5 Antibody Picoband® catalog # A03085. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P55064

### Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Aquaporin 5, which shares 71.7% and 73% amino acid (aa) sequence identity with mouse and rat Aquaporin 5, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml

## Anti-Aquaporin 5/AQP5 Antibody Picoband® (A03085) Images




Western blot analysis of Aquaporin 5 using anti-Aquaporin 5 antibody (A03085). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat lung tissue lysates, Lane 2: mouse lung tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Aquaporin 5 antigen affinity purified polyclonal antibody (Catalog # A03085) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Aquaporin 5 at approximately 28KD. The expected band size for Aquaporin 5 is at 28KD.




Western blot analysis of Aquaporin 5 using anti-Aquaporin 5 antibody (A03085). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: Normal group-rat colon tissue lysates, Lane 2: Model group-rat colon tissue lysates, Lane 3: Traditional Chinese medicine treatment (low concentration)-rat colon tissue lysates, Lane 4: Traditional Chinese medicine treatment (medium concentration)-rat colon tissue lysates, Lane 5: Traditional Chinese medicine treatment (High concentration)-rat colon tissue lysates, Lane 6: Western medicine treatment-rat colon tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Aquaporin 5 antigen affinity purified polyclonal antibody (Catalog # A03085) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with ChemiDoc MP system. The expected band size for Aquaporin 5 is at 28KD.


Western blot analysis of Aquaporin 5 using anti-Aquaporin 5 antibody (A03085). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: MADB106 cells under normal culture conditions, Lane 2: MADB106 cells treated with agonist, Lane 3: MADB106 cells



treated with inhibitor, After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Aquaporin 5 antigen affinity purified polyclonal antibody (Catalog # A03085) at 1:2000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with ChemiDoc MP system. A specific band was detected for Aquaporin 5 at approximately 23KD. The expected band size for Aquaporin 5 is at 28KD.



IHC analysis of Aquaporin 5 using anti-Aquaporin 5 antibody (A03085). Aquaporin 5 was detected in paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Aquaporin 5 Antibody (A03085) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



IHC analysis of Aquaporin 5 using anti-Aquaporin 5 antibody (A03085). Aquaporin 5 was detected in paraffin-embedded section of rat lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Aquaporin 5 Antibody (A03085) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

## 1 Publications Citing This Product

1. PubMed ID: 26668595, Hyperosmotic stress induces cisplatin sensitivity in ovarian cancer cells by stimulating aquaporin-5 expression

Visit [bosterbio.com/anti-aquaporin-5-picoband-trade-antibody-a03085-boster.html](http://bosterbio.com/anti-aquaporin-5-picoband-trade-antibody-a03085-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



For Research Use Only. Not for use in diagnostic procedures.