

Anti-CaMKII alpha/CAMK2A Antibody Picoband®

Catalog Number: A03241-1

About CAMK2A

MAX (Max protein), also called Myc-associated factor x, is the most conserved dimerization component of the MYC-MAX-MXD1 network of basic helix-loop-helix leucine zipper (bHLHZ) transcription factors that regulate cell proliferation, differentiation, and apoptosis. The conservation of the MAX sequence is particularly high in the bHLHZ domain, which is involved in protein-protein interactions and DNA binding. The MAX gene is located on chromosome 14q23 by fluorescence in situ chromosomal hybridization. Both quasisymmetric heterodimers resemble the symmetric MAX homodimer, albeit with marked structural differences in the coiled-coil leucine zipper regions that explain preferential homo- and heteromeric dimerization of these 3 evolutionarily related DNA-binding proteins. MAX acts as a classic tumor suppressor gene. Normal lymphocytes from patients showed absence of methylation of the MAX promoter and biallelic expression of MAX, which ruled out an imprinting-mediated effect on MAX expression. The ability of these cells to divide, differentiate, and apoptose in the absence of Max demonstrated for the first time that these processes can occur via Max- and possibly Myc-independent mechanisms.

Overview


Product Name	Anti-CaMKII alpha/CAMK2A Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CaMKII alpha/CAMK2A Antibody Picoband® catalog # A03241-1. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UQM7

Technical Details

Immunogen	E. coli-derived human CaMKII alpha recombinant protein (Position: L392-R457).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml ELISA, 0.1-0.5ug/ml

Anti-CaMKII alpha/CAMK2A Antibody Picoband® (A03241-1) Images



Western blot analysis of CAMK2A using anti-CAMK2A antibody (A03241-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CAMK2A antigen affinity purified polyclonal antibody (A03241-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for CAMK2A at approximately 54, 60 kDa. The expected band size for CAMK2A is at 54 kDa.

5 Publications Citing This Product

1. PubMed ID: -, Meng, X.,Fu, M.,Wang, S.,Chen, W.,Wang, J.,& Zhang, N.(2021). Naringin ameliorates memory deficits and exerts neuroprotective effects in a mouse model of Alzheimer's disease by regulating multiple metabolic pathways. *Molecular Medicine Reports*,23,332.<https://doi.org/10.3892/mmr.2021.11971>
2. PubMed ID: 27451410, Inhibition of CaMKII α in the central nucleus of amygdala attenuates fentanyl-induced hyperalgesia in rats
3. PubMed ID: 20052459, Sun Z, Niu R, Su K, Wang B, Wang J, Zhang J, Wang J. *Arch Toxicol*. 2010 May;84(5):353-61. Doi: 10.1007/S00204-009-0508-X. Epub 2010 Jan 6. Effects Of Sodium Fluoride On Hyperactivation And Ca²⁺ Signaling Pathway In Sperm From Mice: An In Vivo Study.

Visit bosterbio.com/anti-camkii-alpha-antibody-a03241-1-boster.html to see all 5 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CaMKII alpha/CAMK2A Antibody

For Research Use Only. Not for use in diagnostic procedures.