

Anti-CaMKII alpha/CAMK2A Antibody Picoband® Fluoro550 Conjugated

Catalog Number: A03241-1-Fluoro550

About CAMK2A

MAX (Max protein), also called Myc-associated factor x, is the most conserved dimerization component of the MYC-MAX-MXD1 network of basic helix-loop-helix leucine zipper (bHLHZ) transcription factors that regulate cell proliferation, differentiation, and apoptosis. The conservation of the MAX sequence is particularly high in the bHLHZ domain, which is involved in protein-protein interactions and DNA binding. The MAX gene is located on chromosome 14q23 by fluorescence in situ chromosomal hybridization. Both quasisymmetric heterodimers resemble the symmetric MAX homodimer, albeit with marked structural differences in the coiled-coil leucine zipper regions that explain preferential homo- and heteromeric dimerization of these 3 evolutionarily related DNA-binding proteins. MAX acts as a classic tumor suppressor gene. Normal lymphocytes from patients showed absence of methylation of the MAX promoter and biallelic expression of MAX, which ruled out an imprinting-mediated effect on MAX expression. The ability of these cells to divide, differentiate, and apoptose in the absence of Max demonstrated for the first time that these processes can occur via Max- and possibly Myc-independent mechanisms.

Overview

Product Name	Anti-CaMKII alpha/CAMK2A Antibody Picoband® Fluoro550 Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q9UQM7

Technical Details

Immunogen	E. coli-derived human CaMKII alpha recombinant protein (Position: L392-R457).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	Fluoro550 Excitation Wavelength: 562 nm Emission Wavelength: 576 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

5 Publications Citing This Product

1. PubMed ID: -, Meng, X.,Fu, M.,Wang, S.,Chen, W.,Wang, J.,& Zhang, N.(2021). Naringin ameliorates memory deficits and exerts neuroprotective effects in a mouse model of Alzheimer's disease by regulating multiple metabolic pathways. Molecular Medicine Reports,23,332.https://doi.org/10.3892/mmr.2021.11971
2. PubMed ID: 27451410, Inhibition of CaMKII α in the central nucleus of amygdala attenuates fentanyl-induced hyperalgesia in rats
3. PubMed ID: 20052459, Sun Z, Niu R, Su K, Wang B, Wang J, Zhang J, Wang J. Arch Toxicol. 2010 May;84(5):353-61. Doi: 10.1007/s00204-009-0508-X. Epub 2010 Jan 6. Effects Of Sodium Fluoride On Hyperactivation And Ca²⁺ Signaling Pathway In Sperm From Mice: An In Vivo Study.

Visit bosterbio.com/anti-camkii-alpha-antibody-a03241-1-boster.html to see all 5 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CaMKII alpha/CAMK2A Antibody - Fluoro550

For Research Use Only. Not for use in diagnostic procedures.