

Anti-TROP2/TACSTD2 Antibody Picoband® FITC Conjugated

Catalog Number: A03569-2-FITC

About TACSTD2

Tumor-associated calcium signal transducer 2, also known as Trop-2 and as epithelial glycoprotein-1 antigen (EGP-1), is a protein that in humans is encoded by the TACSTD2 gene. This intronless gene encodes a carcinoma-associated antigen. This antigen is a cell surface receptor that transduces calcium signals. Mutations of this gene have been associated with gelatinous drop-like corneal dystrophy.

Overview

Product Name	Anti-TROP2/TACSTD2 Antibody Picoband® FITC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (Flow Cytometry, IF, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P09758

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human TROP2/TACSTD2, which shares 65% amino acid (aa) sequence identity with mouse and rat TACSTD2.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TROP2/TACSTD2 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.