

Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody

Catalog Number: A03622

About ADRA2B

Alpha-2 adrenergic receptors mediate the catecholamine-induced inhibition of adenylate cyclase through the action of G proteins. The rank order of potency for agonists of this receptor is clonidine > norepinephrine > epinephrine = oxymetazoline > dopamine > p-tyramine = phenylephrine > serotonin > p-synephrine / p-octopamine. For antagonists, the rank order is yohimbine > chlorpromazine > phentolamine > mianserine > spiperone > prazosin > alprenolol > propanolol > pindolol.

Lomasney J.W., Proc. Natl. Acad. Sci. U.S.A. 87:5094-5098(1990). Weinshank R.L., Mol. Pharmacol. 38:681-688(1990). Small K.M., J. Biol. Chem. 276:4917-4922(2001).

Overview

Product Name	Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody
Reactive Species	Human
Description	Boster Bio Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody (Catalog # A03622). Tested in IF applications. This antibody reacts with Human.
Application	IF
Clonality	Polyclonal
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P18089

Technical Details

Immunogen	Synthesized peptide derived from internal of human Adrenergic Receptor alpha-2B.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	lgG
Form	Liquid
Concentration	1 mg/ml




BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitopespecific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunofluorescence: 1:100~1:500



Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody (A03622) Images



Immunofluorescence analysis of HepG2 cells, using Adrenergic Receptor alpha-2B antibody A03622.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody