

Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody

Catalog Number: A03622

About ADRA2B

Alpha-2 adrenergic receptors mediate the catecholamine-induced inhibition of adenylate cyclase through the action of G proteins. The rank order of potency for agonists of this receptor is clonidine > norepinephrine > epinephrine = oxymetazoline > dopamine > p-tyramine = phenylephrine > serotonin > p-synephrine / p-octopamine. For antagonists, the rank order is yohimbine > chlorpromazine > phentolamine > mianserine > spiperone > prazosin > alprenolol > propanolol > pindolol.

Lomasney J.W., Proc. Natl. Acad. Sci. U.S.A. 87:5094-5098(1990).

Weinshank R.L., Mol. Pharmacol. 38:681-688(1990).

Small K.M., J. Biol. Chem. 276:4917-4922(2001).

Overview


Product Name	Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody
Reactive Species	Human
Description	Boster Bio Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody (Catalog # A03622). Tested in IF applications. This antibody reacts with Human.
Application	IF
Clonality	Polyclonal
Formulation	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P18089

Technical Details

Immunogen	Synthesized peptide derived from internal of human Adrenergic Receptor alpha-2B.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunofluorescence: 1:100~1:500

Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody (A03622) Images



Immunofluorescence analysis of HepG2 cells, using Adrenergic Receptor alpha-2B antibody A03622.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Adrenergic Receptor alpha-2B ADRA2B Antibody