

## Anti-Fanca/FAA Antibody Picoband®

Catalog Number: A03662-2

### About Fanca

Fanconi anaemia, complementation group A, also known as FAA, FACA and FANCA, is a protein which in humans is encoded by the FANCA gene. Acts upstream of or within several processes, including gonad development; regulation of CD40 signaling pathway; and regulation of regulatory T cell differentiation. Predicted to be located in nucleoplasm. Predicted to be part of Fanconi anaemia nuclear complex. Is expressed in several structures, including gut; liver; lung; reproductive system; and spleen. Used to study Fanconi anemia complementation group A. Human ortholog(s) of this gene implicated in several diseases, including Fanconi anemia (multiple); acute myeloid leukemia; esophagus squamous cell carcinoma; ovarian disease (multiple); and skin melanoma. Orthologous to human FANCA (FA complementation group A).

### Overview


Product Name	Anti-Fanca/FAA Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-Fanca/FAA Antibody Picoband® catalog # A03662-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9JL70

### Technical Details


Immunogen	E.coli-derived mouse Fanca/FAA recombinant protein (Position: E1272-F1439).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse, Rat Flow Cytometry (Fixed), 1-3 ug/1x10 <sup>6</sup> cells, Mouse ELISA, 0.1-0.5 ug/ml, -

## Anti-Fanca/FAA Antibody Picoband® (A03662-2) Images



Western blot analysis of Fanca/FAA using anti-Fanca/FAA antibody (A03662-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat testis tissue lysates, Lane 2: mouse thymus tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Fanca/FAA antigen affinity purified polyclonal antibody (Catalog # A03662-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Fanca/FAA at approximately 180 kDa. The expected band size for Fanca/FAA is at 161 kDa.



Flow Cytometry analysis of ANA-1 cells using anti-Fanca/FAA antibody (A03662-2). Overlay histogram showing ANA-1 cells stained with A03662-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Fanca/FAA Antibody (A03662-2, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Fanca/FAA Antibody

For Research Use Only. Not for use in diagnostic procedures.