

Anti-KChIP3 Antibody

Catalog Number: A03700-1

About KCNIP3

CAND1 is also known as TIP120A, and TATA-binding protein-interacting protein 120A. The SCF complex consists of the invariable components Skp1, Cul1, and Rbx1 as well as a variable F-box protein, and functions as an E3 ubiquitin ligase. E3 ubiquitin ligases regulate various physiological processes. CAND1 binds to Cul1 and potentially regulates the SCF complex. CAND1 physically associates with Cul1 in the nucleus and this interaction is mediated by a central region of Cul1 distinct from its binding sites for Skp1 and Rbx1. CAND1 selectively binds to unneddylated CUL1 and is dissociated by CUL1 neddylation. CAND1 forms a ternary complex with CUL1 and ROC1.

Overview

| Product Name | Anti-KChIP3 Antibody |
|----------------------|--|
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-KChIP3 Antibody catalog # A03700-1. Tested in IHC applications. This antibody reacts with Human, Mouse, Rat. |
| Application | IHC |
| Clonality | Polyclonal 5A6 |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q9Y2W7 |

Technical Details

| Immunogen | Synthetic Peptide |
|----------------------------|--|
| Predicted Reactive Species | Canine, Equine, Guinea Pig, Rabbit |
| Cross Reactivity | No cross reactivity with other proteins. |
| Isotype | lgG |
| Form | Liquid |
| Concentration | This antibody's concentration is >0.5mg/ml. |
| Purification | Immunogen affinity purified |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this |




BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com


| | kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: IHC, 1:100-200 |
|--|---|
|--|---|



Anti-KChIP3 Antibody (A03700-1) Images



Immunohistochemistry (IHC) analysis of paraffin-embedded Rat Brain Tissue using KChIP3 Rabbit Polyclonal antibody diluted at 1:200.



Immunohistochemistry (IHC) analysis of paraffin-embedded Mouse Brain Tissue using KChIP3 Rabbit Polyclonal antibody diluted at 1:200.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-KChIP3 Antibody