BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Histamine N-methyltransferase HNMT Antibody

Catalog Number: A03807

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview


Product Name	Anti-Histamine N-methyltransferase HNMT Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Histamine N-methyltransferase HNMT Antibody catalog # A03807. Tested in WB,IP applications. This antibody reacts with Human,Mouse.
Application	IP, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P50135

Technical Details


Immunogen	Recombinant fusion protein of human HNMT(NP_008826.1).
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	WB: 1:500-1:2000 IP: 1:20-1:50 RRM2B/p53R2 antibody is tested in Western Blot and ELISA and suitable for IP, IHC, and IF. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 41 kDa in size corresponding to RRM2B1 by western blotting in the appropriate cell lysate or extract.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Histamine N-methyltransferase HNMT Antibody (A03807) Images



Western blot analysis of extracts of various cell lines, using HNMT antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit .Exposure time: 30s.



Immunoprecipitation analysis of 200ug extracts of HT-29 cells using 1ug HNMT antibody . Western blot was performed from the immunoprecipitate using HNMT antibody at a dilution of 1:1000.

Anti-Histamine N-methyltransferase HNMT Antibody

For Research Use Only. Not for use in diagnostic procedures.