

## Anti-Claudin 7/CLDN-7 Antibody Picoband™

Catalog Number: A03851-3

### About CLDN7

Claudin-7 is a protein that in humans is encoded by the CLDN7 gene. This gene encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This gene is expressed constitutively in the mammary epithelium throughout development, and might be involved in vesicle trafficking to the basolateral membrane. It is essential for NaCl homeostasis in distal nephrons. The knockout mice lacking this gene showed severe salt wasting, chronic dehydration, and growth retardation, and died within 12 days after birth. Alternatively spliced transcript variants have been found for this gene.

### Overview

Product Name	Anti-Claudin 7/CLDN-7 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Claudin 7/CLDN-7 Antibody Picoband™ catalog # A03851-3. Tested in ELISA, Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O95471

### Technical Details

Immunogen	E.coli-derived human Claudin 7/CLDN-7 recombinant protein (Position: F92-V211).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.25 ug/ml, Human</p> <p>Immunohistochemistry (Paraffin-embedded Section), 1-2ug/ml, Human</p> <p>Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human</p> <p>Immunofluorescence, 5 ug/ml, Human</p> <p>Flow Cytometry, 1-3ug/1x10<sup>6</sup> cells, Human</p> <p>Direct ELISA, 0.1-0.5ug/ml, Human</p>

## Anti-Claudin 7/CLDN-7 Antibody Picoband™ (A03851-3) Images




Figure 1. Western blot analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Caco-2 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Claudin 7/CLDN-7 antigen affinity purified polyclonal antibody (Catalog # A03851-3) at 0.25 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Claudin 7/CLDN-7 at approximately 22 kDa. The expected band size for Claudin 7/CLDN-7 is at 22 kDa.




Figure 2. IHC analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3).

Claudin 7/CLDN-7 was detected in a paraffin-embedded section of human renal clear cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.





Figure 3. IHC analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3).

Claudin 7/CLDN-7 was detected in a paraffin-embedded section of human gallbladder adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Figure 4. IHC analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3).



Claudin 7/CLDN-7 was detected in a paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 5. IF analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3).

Claudin 7/CLDN-7 was detected in a paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3) overnight at 4°C. Biotin conjugated goat anti-rabbit IgG (BA1003) was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using DyLight®488 Conjugated Avidin (BA1128). The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. IF analysis of Claudin 7/CLDN-7 using anti-Claudin 7/CLDN-7 antibody (A03851-3).

Claudin 7/CLDN-7 was detected in an immunocytochemical section of MCF-7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 7. Flow Cytometry analysis of Caco-2 cells using anti-Claudin 7/CLDN-7 antibody (A03851-3).

Overlay histogram showing Caco-2 cells stained with A03851-3 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Claudin 7/CLDN-7 Antibody (A03851-3, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



## Anti-Claudin 7/CLDN-7 Antibody