

## Anti-STOML2 Antibody Picoband®

Catalog Number: A04108-1

### About STOML2

Enables GTPase binding activity; T cell receptor binding activity; and cardiolipin binding activity. Involved in intracellular calcium ion homeostasis; mitochondrion organization; and protein complex oligomerization. Acts upstream of or within T cell receptor signaling pathway. Located in several cellular components, including immunological synapse; membrane raft; and mitochondrial envelope.

### Overview


Product Name	Anti-STOML2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-STOML2 Antibody Picoband® catalog # A04108-1. Tested in WB, IHC, IP, Flow Cytometry, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IP, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9UJZ1

### Technical Details


Immunogen	E.coli-derived human STOML2 recombinant protein (Position: Q43-S293). Human STOML2 shares 98.8% amino acid (aa) sequence identity with both mouse and rat STOML2.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Human Immunoprecipitation, 0.5-2 ug/ml, Human Flow Cytometry (Fixed), 1-3 ug/1x10 <sup>6</sup> cells, Human ELISA, 0.1-0.5 ug/ml




## Anti-STOML2 Antibody Picoband® (A04108-1) Images




Western blot analysis of STOML2 using anti-STOML2 antibody (A04108-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human 293T whole cell lysates, Lane 2: human HeLa whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human MCF-7 whole cell lysates, Lane 5: rat brain tissue lysates, Lane 6: rat C6 whole cell lysates, Lane 7: mouse brain tissue lysates, Lane 8: mouse NIH/3T3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-STOML2 antigen affinity purified polyclonal antibody (A04108-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for STOML2 at approximately 39 kDa. The expected band size for STOML2 is at 39 kDa.



IHC analysis of STOML2 using anti-STOML2 antibody (A04108-1). STOML2 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-STOML2 Antibody (A04108-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Immunoprecipitating STOML2 in jurkat whole cell lysate. Western blot analysis of STOML2 using anti-STOML2 antibody (A04108-1). Lane 1: jurkat whole cell lysates (30ug), Lane 2: Rabbit control IgG instead of anti-STOML2 antibody in jurkat whole cell lysate, Lane 3: anti-STOML2 antibody (2ug) + jurkat whole cell lysate (500ug). After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-STOML2 antigen affinity purified polyclonal antibody (A04108-1) at a dilution of 0.5 ug/mL and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for STOML2 at approximately 39 kDa. The expected band size for STOML2 is at 39 kDa.



Flow Cytometry analysis of Jurkat cells using anti-STOML2 antibody (A04108-1). Overlay histogram showing Jurkat cells stained with A04108-1 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-STOML2 Antibody (A04108-1, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-STOML2 Antibody

For Research Use Only. Not for use in diagnostic procedures.