BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Protein 4.1 EPB41 Antibody

Catalog Number: A04124

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Protein 4.1 EPB41 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Protein 4.1 EPB41 Antibody catalog # A04124. Tested in WB,ELISA,IHC,IF,ICC applications. This antibody reacts with Human,Mouse,Rat.
Application	ELISA, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P11171

Technical Details


Immunogen	Recombinant full length Human EPB41.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	WB: 1:500-1:2000 ELISA: 1:5000-1:10000 IHC: 1:100-1:500 IF: 1:100-1:500

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

	ICC: 1:100-1:500 Affinity purified Bach-2 antibody has been tested for use in ELISA, Immunoprecipitation, and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 91-140 kDa in size corresponding to Bach2 protein by western blotting in the appropriate cell lysate or extract.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Protein 4.1 EPB41 Antibody (A04124) Images



Tissue/cell: rat spleen tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-EPB41 Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining

Anti-Protein 4.1 EPB41 Antibody

For Research Use Only. Not for use in diagnostic procedures.