

Anti-NAGA Antibody Picoband® FITC Conjugated

Catalog Number: A04163-1-FITC

About NAGA

NAGA encodes the lysosomal enzyme alpha-N-acetylgalactosaminidase, which cleaves alpha-N-acetylgalactosaminyl moieties from glycoconjugates. Mutations in NAGA have been identified as the cause of Schindler disease types I and II (type II also known as Kanzaki disease).

Overview

Product Name	Anti-NAGA Antibody Picoband® FITC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P17050

Technical Details

Immunogen	E.coli-derived human NAGA recombinant protein (Position: D43-R329). Human NAGA shares 86.7% and 85.4% amino acid (aa) sequence identity with mouse and rat NAGA, respectively.
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your



reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-NAGA Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.