BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Probable hydrolase PNKD PNKD Antibody

Catalog Number: A04242

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Probable hydrolase PNKD PNKD Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Probable hydrolase PNKD PNKD Antibody catalog # A04242. Tested in WB,ICC/IF applications. This antibody reacts with Human,Mouse.
Application	IF, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8N490


Technical Details

Immunogen	Recombinant fusion protein of human PNKD(NP_056303.3).
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	WB: 1:500-1:2000 ICC/IF: 1:50-1:200 Affinity purified Bach-2 antibody has been tested for use in ELISA, Immunoprecipitation, and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 91-140 kDa in size corresponding to Bach2 protein by western blotting in the appropriate cell lysate or extract.


BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Probable hydrolase PNKD PNKD Antibody (A04242) Images



Western blot analysis of extracts of various cell lines, using PNKD antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit .Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using PNKD antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Anti-Probable hydrolase PNKD PNKD Antibody

For Research Use Only. Not for use in diagnostic procedures.