

## Anti-PI4KB Antibody Picoband®

Catalog Number: A04249-2

### About PI4KB

Phosphatidylinositol 4-kinase beta is an enzyme that in humans is encoded by the PI4KB gene. Inositol phospholipids have an important role in intracellular signaling in response to hormones, growth factors and neurotransmitters. Phosphatidylinositol 4-kinase phosphorylates phosphatidylinositol (PI) to phosphatidylinositol-4-phosphate (PIP). In a second step, PIP is further phosphorylated to phosphatidylinositol-4,5-bisphosphate (PIP2), and PIP2 is subsequently hydrolyzed by phospholipase C, producing the two intracellular second messengers, inositol 1,4,5-triphosphate (IP3) and diacylglycerol (DAG). PI4K230, PI4K92 and PI4K55 are three PI4 kinase isoforms that have been characterized and classified according to their molecular weights of 230, 92 and 55 kD. Previously, PI4 kinases were classified into type II and III enzymes. All isoforms are located on distinct membranes and cellular compartments suggesting various tasks. PI4K230 is located at the endoplasmic reticulum and outer membranes of mitochondria, PI4K92 at the Golgi apparatus and endoplasmic reticulum, and PI4K55 at the plasma membrane and endosomes. PI4K230 is predominantly expressed in brain and moderately sensitive to wortmannin as well as specifically and irreversibly inhibited by cyclitol derivatives.

### Overview


Product Name	Anti-PI4KB Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PI4KB Antibody Picoband® catalog # A04249-2. Tested in ELISA, IHC, WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9UBF8

### Technical Details


Immunogen	E.coli-derived human PI4KB recombinant protein (Position: D375-D671). Human PI4KB shares 100% and 99.7% amino acid (aa) sequence identity with mouse and rat PI4KB, respectively
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).

Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Human Flow Cytometry (Fixed), 1-3 ug/1x10 <sup>6</sup> cells, Human ELISA, 0.1-0.5 ug/ml, -


## Anti-PI4KB Antibody Picoband® (A04249-2) Images



Western blot analysis of PI4KB using anti-PI4KB antibody (A04249-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: human Hela whole cell lysates, Lane 3: rat brain tissue lysates, Lane 4: mouse brain tissue lysates, Lane 5: mouse lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PI4KB antigen affinity purified polyclonal antibody (Catalog # A04249-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PI4KB at approximately 95 kDa. The expected band size for PI4KB is at 91,91-110 kDa.




IHC analysis of PI4KB using anti-PI4KB antibody (A04249-2). PI4KB was detected in a paraffin-embedded section of human skeletal muscle tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-PI4KB Antibody (A04249-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Flow Cytometry analysis of JK cells using anti-PI4KB antibody (A04249-2). Overlay histogram showing JK cells stained with A04249-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PI4KB Antibody (A04249-2, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Flow Cytometry analysis of U251 cells using anti-PI4KB antibody (A04249-2). Overlay histogram showing U251 cells stained with A04249-2 (Blue line). To facilitate intracellular



staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PI4KB Antibody (A04249-2, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

## Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-PI4KB Antibody

For Research Use Only. Not for use in diagnostic procedures.