

Anti-VIP Receptor 1/VIPR1 Antibody Picoband®

Catalog Number: A04345

About VIPR1

VIPR1 (Vasoactive intestinal polypeptide receptor 1), also known as VIPR, HVR1, is a protein that in humans is encoded by the VIPR1 gene. Distinct subsets of neural, respiratory, gastrointestinal, and immune cells bear specific high-affinity G protein-coupled receptors for VIP, such as VIPR1. The VIPR1 gene is mapped on 3p22.1. The VIPR1 gene was found to span approximately 22 kb and to be comprised of 13 exons (ranging from 42 to 1,400 bp) and 12 introns (ranging from 0.3 to 6.1 kb). One encodes a VIP receptor consisting of 460 amino acids and having 7 putative transmembrane domains, as do other G protein-coupled receptors. Patients with idiopathic achalasia show a significant difference in the distribution of SNPs affecting VIPR1.

Overview


Product Name	Anti-VIP Receptor 1/VIPR1 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-VIP Receptor 1/VIPR1 Antibody Picoband® catalog # A04345. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P32241

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human VIP Receptor 1, which shares 84% amino acid (aa) sequence identity with both mouse and rat VIP Receptor 1.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml

Anti-VIP Receptor 1/VIPR1 Antibody Picoband® (A04345) Images



Western blot analysis of VIP Receptor 1 using anti-VIP Receptor 1 antibody (A04345). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human A431 cell lysates, Lane 2: human PANC-1 cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VIP Receptor 1 antigen affinity purified polyclonal antibody (Catalog # A04345) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for VIP Receptor 1 at approximately 52KD. The expected band size for VIP Receptor 1 is at 52KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-VIP Receptor 1/VIPR1 Antibody

For Research Use Only. Not for use in diagnostic procedures.