

Anti-Vasohibin VASH1 Antibody

Catalog Number: A04402-1

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview

Product Name	Anti-Vasohibin VASH1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Vasohibin VASH1 Antibody catalog # A04402-1. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% stabilizing protein and 0.02% sodium azide. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q7L8A9


Technical Details

Immunogen	The antiserum was produced against synthesized peptide derived from human VASH1. AA range:261-310
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	Immunogen affinity purified


Suggested Dilutions

WB 1:500-1:2000
ELISA 1:40000


Anti-Vasohibin VASH1 Antibody (A04402-1) Images




Western Blot analysis of various cells using Vasohibin Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western blot analysis of VASH1 Antibody. The lane on the right is blocked with the VASH1 peptide.



Immunohistochemical analysis of paraffin-embedded human colon. 1, primary Antibody was diluted at 1:200 (4° overnight). 2, EDTA pH 9.0 was used for antigen retrieval. 3 Reday to use Secondary antibody was used at 37°C, 30min



Immunohistochemical analysis of paraffin-embedded human brain. 1, primary Antibody was diluted at 1:200 (4° overnight). 2, EDTA pH 9.0 was used for antigen retrieval 3 Reday to use Secondary antibody was used at 37°C, 30min

[Anti-Vasohibin VASH1 Antibody](#)

For Research Use Only. Not for use in diagnostic procedures.