BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Vasohibin VASH1 Antibody

Catalog Number: A04402-1

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Vasohibin VASH1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Vasohibin VASH1 Antibody catalog # A04402-1. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% stabilizing protein and 0.02% sodium azide. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q7L8A9

Technical Details


Immunogen	The antiserum was produced against synthesized peptide derived from human VASH1. AA range:261-310
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	Immunogen affinity purified
Suggested Dilutions	WB 1:500-1:2000

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


	ELISA 1:40000 Anti-Thymidylate Synthase has been tested by western blot and is suitable for use in ELISA, immunoprecipitation, immunofluorescence microscopy, immunohistochemistry and immunoblotting. The antibody recognizes the expected additional band corresponding to the ternary complex of hTS-dFUMP-reduced folate in HeLa cells treated with the TS inhibitor 5-FUdR. This event occurs in most human breast, colorectal, gastric, head and neck carcinomas. The antibody recognizes the 36,000 MW hTS. Reactivity in other immunoassays is unknown.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-Vasohibin VASH1 Antibody (A04402-1) Images




Western Blot analysis of various cells using Vasohibin Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western blot analysis of VASH1 Antibody. The lane on the right is blocked with the VASH1 peptide.



Immunohistochemical analysis of paraffin-embedded human colon. 1, primary Antibody was diluted at 1:200 (4° overnight). 2, EDTA pH 9.0 was used for antigen retrieval. 3 Reday to use Secondary antibody was used at 37°C, 30min



Immunohistochemical analysis of paraffin-embedded human brain. 1, primary Antibody was diluted at 1:200 (4° overnight). 2, EDTA pH 9.0 was used for antigen retrieval 3 Reday to use Secondary antibody was used at 37°C, 30min

For Research Use Only. Not for use in diagnostic procedures.