

Anti-GIT2 Antibody

Catalog Number: A04441

About GIT2

Thymidylate synthase catalyzes the methylation of deoxyuridylate to deoxythymidylate using 5,10-methylenetetrahydrofolate (methylene-THF) as a cofactor. This function maintains the dTMP (thymidine-5-prime monophosphate) pool critical for DNA replication and repair. The enzyme has been of interest as a target for cancer chemotherapeutic agents. It is considered to be the primary site of action for 5-fluorouracil, 5-fluoro-2-prime-deoxyuridine, and some folate analogs. Expression of this gene and that of a naturally occurring antisense transcript rTS alpha vary inversely when cell-growth progresses from late-log to plateau phase. Diseases associated with Thymidylate synthase include Rectal Neoplasm and Dihydropyrimidine Dehydrogenase Deficiency. Anti-Thymidylate synthase is useful for researchers interested in Circadian Rhythm, Metabolism and cell cycle research.

Overview

Product Name	Anti-GIT2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-GIT2 Antibody catalog # A04441. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal 3F8
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q14161

Technical Details

Immunogen	Synthesized peptide derived from the Internal region of human GIT2. at AA range: 330-410
Predicted Reactive Species	Bovine, Canine, Equine, Goat, Guinea Pig, Pig, Rabbit, Sheep
Isotype	IgG
Form	Liquid
Concentration	1 mg/mL.
Purification	Immunogen affinity purified

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.


If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

WB, 1:500-1:2000

Anti-GIT2 Antibody (A04441) Images



Western Blot (WB) analysis of HepG2 cells using GIT2 Polyclonal antibody.



Western Blot (WB) analysis of specific cells using GIT2 Polyclonal antibody.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-GIT2 Antibody