

Anti-Noggin Antibody Picoband® Fluoro594 Conjugated

Catalog Number: A04448-1-Fluoro594

About NOG

The secreted polypeptide, encoded by NOG gene, binds and inactivates members of the transforming growth factor-beta (TGF-beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP4). By diffusing through extracellular matrices more efficiently than members of the TGF-beta superfamily, this protein may have a principal role in creating morphogenic gradients. The protein appears to have pleiotropic effect, both early in development as well as in later stages. It was originally isolated from *Xenopus* based on its ability to restore normal dorsal-ventral body axis in embryos that had been artificially ventralized by UV treatment. The results of the mouse knockout of the ortholog suggest that it is involved in numerous developmental processes, such as neural tube fusion and joint formation. Recently, several dominant human NOG mutations in unrelated families with proximal symphalangism (SYM1) and multiple synostoses syndrome (SYNS1) were identified; both SYM1 and SYNS1 have multiple joint fusion as their principal feature, and map to the same region (17q22) as this gene. All of these mutations altered evolutionarily conserved amino acid residues. The amino acid sequence of this human gene is highly homologous to that of *Xenopus*, rat and mouse.

Overview

Product Name	Anti-Noggin Antibody Picoband® Fluoro594 Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q13253

Technical Details

Immunogen	E. coli-derived human Noggin recombinant protein (Position: L43-T153).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	Fluoro594 Excitation Wavelength: 593 nm Emission Wavelength: 618 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

1 Publications Citing This Product

1. PubMed ID: 21573506, Expression of NANOG in human gliomas and its relationship with undifferentiated glioma cells

Visit bosterbio.com/anti-noggin-picoband-trade-antibody-a04448-1-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Noggin Antibody - Fluoro594

For Research Use Only. Not for use in diagnostic procedures.