

Anti-Agrin/AGRN Picoband® Antibody Cy3 Conjugated

Catalog Number: A04649-Cy3

About AGRN

Agrin is a protein that in humans is encoded by the AGRN gene. It is mapped to 1p36.33. This gene encodes one of several proteins that are critical in the development of the neuromuscular junction (NMJ), as identified in mouse knock-out studies. The encoded protein contains several laminin G, Kazal type serine protease inhibitor, and epidermal growth factor domains. Additional post-translational modifications occur to add glycosaminoglycans and disulfide bonds. In one family with congenital myasthenic syndrome affecting limb-girdle muscles, a mutation in this gene was found. Alternative splicing results in multiple transcript variants encoding different isoforms.

Overview

Product Name	Anti-Agrin/AGRN Picoband® Antibody Cy3 Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, IHC, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	O00468

Technical Details

Immunogen	E.coli-derived human Agrin/AGRN recombinant protein (Position: S1864-P2068).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Cy3 Excitation Wavelength: 554 nm Emission Wavelength: 568 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Agrin/AGRN Antibody - Cy3

For Research Use Only. Not for use in diagnostic procedures.