

## Anti-PUMA/BBC3 Antibody Picoband®

Catalog Number: A04899-3

### About BBC3

The p53 upregulated modulator of apoptosis, or PUMA, is a pro-apoptotic member of the Bcl-2 protein family. The PUMA gene is located at 19q. PUMA transcript is contained within 4 exons, with the presumptive initiation codon in exon 2. The predicted 193-amino acid PUMA protein shares 91% amino acid identity with the murine sequence. Bcl-2 family members can form hetero- or homodimers, and they act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The expression of PUMA is regulated by the tumor suppressor p53, and PUMA has been shown to be involved in p53-mediated apoptosis. Additionally, PUMA encodes 2 BH3 domain-containing proteins, PUMA-alpha and PUMA-beta, that are produced through the use of an alternative first exon and are induced in cells following p53 activation. Furthermore, PUMA couples the nuclear and cytoplasmic proapoptotic functions of p53.

### Overview


Product Name	Anti-PUMA/BBC3 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-PUMA/BBC3 Antibody Picoband® catalog # A04899-3. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9BXH1

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human PUMA/BBC3, which shares 95.7% amino acid (aa) sequence identity with mouse and rat BBC3.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml/ml, Human

## Anti-PUMA/BBC3 Antibody Picoband® (A04899-3) Images



Western blot analysis of PUMA/BBC3 using anti-PUMA/BBC3 antibody (A04899-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human K562 whole cell lysates, Lane 3: human Hela whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PUMA/BBC3 antigen affinity purified polyclonal antibody (Catalog # A04899-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PUMA/BBC3 at approximately 18 kDa. The expected band size for PUMA/BBC3 is at 18 kDa.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-PUMA/BBC3 Antibody

For Research Use Only. Not for use in diagnostic procedures.