

Anti-Flt3 ligand/Flt3lg Antibody Picoband®

Catalog Number: A04987-1

About Flt3lg

FLT3LG (FMS-Related Tyrosine Kinase 3 Ligand) also called FLT3 LIGAND, FL or FLT3L, is a protein which in humans is encoded by the FLT3LG gene. FLT3LG controls the development of DCs and is particularly important for plasmacytoid DCs and CD8-positive classical DCs and their CD103-positive tissue counterparts. Flt3 ligand (FL) is a hematopoietic four helical bundle cytokine. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors. Lyman et al. (1993) found that Flt3 ligand stimulated proliferation of hematopoietic progenitor cells isolated from mouse fetal liver or adult mouse bone marrow. Hannum et al. (1994) concluded that FL enhances the response of stem and primitive progenitor cells to other growth factors to generate all myeloid lineages except erythroid cells. Assays of C-terminally truncated FL proteins confirmed that the N-terminal half conferred FL activity. Depletion of the Pi3k -Mtor negative regulator Pten facilitated Flt3l-driven DC development in culture.

Overview


Product Name	Anti-Flt3 ligand/Flt3lg Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-Flt3 ligand/Flt3lg Antibody Picoband® catalog # A04987-1. Tested in ELISA, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	A0A0G2K6K8

Technical Details

Immunogen	E. coli-derived rat Flt3 ligand recombinant protein (Position: T28-Q189).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Mouse, Rat ELISA, 0.1-0.5ug/ml, -

Anti-Flt3 ligand/Flt3lg Antibody Picoband® (A04987-1) Images



Western blot analysis of Flt3 ligand using anti-Flt3 ligand antibody (A04987-1). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat thymus tissue lysates, Lane 2: mouse thymus tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Flt3 ligand antigen affinity purified polyclonal antibody (A04987-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for Flt3 ligand at approximately 26 kDa. The expected band size for Flt3 ligand is at 26 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Flt3 ligand/Flt3lg Antibody

For Research Use Only. Not for use in diagnostic procedures.