

Anti-Flt3 ligand/Flt3lg Antibody Picoband® Biotin Conjugated

Catalog Number: A04987-1-Biotin

About Flt3lg

FLT3LG (FMS-Related Tyrosine Kinase 3 Ligand) also called FLT3 LIGAND, FL or FLT3L, is a protein which in humans is encoded by the FLT3LG gene. FLT3LG controls the development of DCs and is particularly important for plasmacytoid DCs and CD8-positive classical DCs and their CD103-positive tissue counterparts. Flt3 ligand (FL) is a hematopoietic four helical bundle cytokine. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors. Lyman et al. (1993) found that Flt3 ligand stimulated proliferation of hematopoietic progenitor cells isolated from mouse fetal liver or adult mouse bone marrow. Hannum et al. (1994) concluded that FL enhances the response of stem and primitive progenitor cells to other growth factors to generate all myeloid lineages except erythroid cells. Assays of C-terminally truncated FL proteins confirmed that the N-terminal half conferred FL activity. Depletion of the Pi3k -Mtor negative regulator Pten facilitated Flt3l-driven DC development in culture.

Overview

| | |
|----------------------|--|
| Product Name | Anti-Flt3 ligand/Flt3lg Antibody Picoband® Biotin Conjugated |
| Reactive Species | Mouse, Rat |
| Application | WB, IHC, ELISA |
| Clonality | Polyclonal |
| Formulation | Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ . |
| Storage Instructions | At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. |
| Host | Rabbit |
| Uniprot ID | A0A0G2K6K8 |

Technical Details

| | |
|------------------|---|
| Immunogen | E. coli-derived rat Flt3 ligand recombinant protein (Position: T28-Q189). |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Purification | Immunogen affinity purified. |
| Conjugate | Biotin |

Suggested Dilutions

Western blot, Optimal dilutions should be determined by end users.
Immunohistochemistry (Paraffin-embedded Section), Optimal dilutions should be determined by end users.
ELISA, Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Flt3 ligand/Flt3Ig Antibody - Biotin

For Research Use Only. Not for use in diagnostic procedures.