

Anti-ACADS/SCAD Antibody Picoband™

Catalog Number: A05028-1

About ACADS

Acyl-CoA dehydrogenase, C-2 to C-3 short chain is an enzyme that in humans is encoded by the ACADS gene. This gene encodes a tetrameric mitochondrial flavoprotein, which is a member of the acyl-CoA dehydrogenase family. This enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation pathway. Mutations in this gene have been associated with short-chain acyl-CoA dehydrogenase (SCAD) deficiency. Alternative splicing results in two variants which encode different isoforms.

Overview

Product Name	Anti-ACADS/SCAD Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ACADS/SCAD Antibody Picoband™ catalog # A05028-1. Tested in IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P16219

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human ACADS/SCAD, identical to the related mouse and rat sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.25-0.5ug/ml, Human, Mouse, Rat

Immunocytochemistry/Immunofluorescence, 5ug/ml, Human

Anti-ACADS/SCAD Antibody Picoband™ (A05028-1) Images




Figure 1. Western blot analysis of ACADS/SCAD using anti-ACADS/SCAD antibody (A05028-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,
Lane 2: human A549 whole cell lysates,
Lane 3: human U87 whole cell lysates,
Lane 4: human HepG2 whole cell lysates,
Lane 5: human Hek293 whole cell lysates,
Lane 6: rat liver tissue lysates,
Lane 7: rat kidney tissue lysates,
Lane 8: mouse liver tissue lysates,
Lane 9: mouse kidney tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ACADS/SCAD antigen affinity purified polyclonal antibody (Catalog # A05028-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ACADS/SCAD at approximately 44KD. The expected band size for ACADS/SCAD is at 44KD.




Figure 2. IF analysis of ACADS/SCAD using anti-ACADS/SCAD antibody (A05028-1).

ACADS/SCAD was detected in immunocytochemical section of MCF-7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-ACADS/SCAD Antibody (A05028-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ACADS/SCAD Antibody