

Anti-CTNNBL1 Antibody Picoband® Cy3 Conjugated

Catalog Number: A05286-1-Cy3

About CTNNBL1

Beta-catenin-like protein 1 is a protein that in humans is encoded by the CTNNBL1 gene. The protein encoded by this gene is a component of the pre-mRNA-processing factor 19-cell division cycle 5-like (PRP19-CDC5L) protein complex, which activates pre-mRNA splicing and is an integral part of the spliceosome. The encoded protein is also a nuclear localization sequence binding protein, and binds to activation-induced deaminase and is important for antibody diversification. This gene may also be associated with the development of obesity. Alternative splicing results in multiple transcript variants. A pseudogene of this gene has been defined on the X chromosome.

Overview

Product Name	Anti-CTNNBL1 Antibody Picoband® Cy3 Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q8WYA6

Technical Details

Immunogen	E.coli-derived human CTNNBL1 recombinant protein (Position: Q122-E231). Human CTNNBL1 shares 97.3% and 98.2% amino acid (aa) sequence identity with mouse and rat CTNNBL1, respectively.
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Cy3 Excitation Wavelength: 554 nm Emission Wavelength: 568 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CTNNB1 Antibody - Cy3

For Research Use Only. Not for use in diagnostic procedures.