

Anti-Nefh Antibody Picoband®

Catalog Number: A05307

About Nefh

Neurofilaments are type IV intermediate filament heteropolymers composed of light, medium, and heavy chains. Neurofilaments comprise the axoskeleton and functionally maintain neuronal caliber. They may also play a role in intracellular transport to axons and dendrites. This gene encodes the heavy neurofilament protein. This protein is commonly used as a biomarker of neuronal damage and susceptibility to amyotrophic lateral sclerosis (ALS) has been associated with mutations in this gene.

Overview


Product Name	Anti-Nefh Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Nefh Antibody Picoband® catalog # A05307. Tested in ELISA, Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P19246

Technical Details


Immunogen	E.coli-derived mouse Nefh recombinant protein (Position: Y109-E466).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.25ug/ml, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Mouse ELISA, 0.1-0.5ug/ml, -


Anti-Nefh Antibody Picoband® (A05307) Images



Western blot analysis of Nefh using anti-Nefh antibody (A05307). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Nefh antigen affinity purified polyclonal antibody (Catalog # A05307) at 0.25 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Nefh at approximately 117-220KD. The expected band size for Nefh is at 117KD.




IHC analysis of Nefh using anti-Nefh antibody (A05307). Nefh was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Nefh Antibody (A05307) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




IHC analysis of Nefh using anti-Nefh antibody (A05307). Nefh was detected in paraffin-embedded section of mouse spinal cord tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Nefh Antibody (A05307) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.


IHC analysis of Nefh using anti-Nefh antibody (A05307). Nefh was detected in a paraffin-embedded section of human cerebellum tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Nefh Antibody (A05307) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as




secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Flow Cytometry analysis of Neuro-2a cells using anti-Nefh antibody (A05307). Overlay histogram showing Neuro-2a cells stained with A05307 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Nefh Antibody (A05307, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.



Images of immunofluorescence staining in each group. (A) Pictures represent the expression of NF-200, GAP-43 and MAP-2, respectively. Rats in each group were sacrificed at Day 14 after the MCAO. Scale bar = 50 um. (B) Immunofluorescence analysis of NF-200 (n = 3), * p < 0.05, as compared with the AAV-empty group, # p < 0.05, as compared with the AAV-NT-1 group. (C) Immunofluorescence analysis of GAP-43 (n = 3), * p < 0.01, as compared with the AAV-empty group, # p < 0.01, as compared with the AAV-NT-1 group. (D) Immunofluorescence analysis of MAP-2 (n = 3), * p < 0.05, as compared with the AAV-empty group. Index in PubMed under a CC BY license. PMID: 29487502



Effects of Netrin-1 overexpression and/or JNK inhibition on the axonal regeneration after the MCAO. (A) The representative images of western-blot analysis for NF-200, GAP-43 and MAP-2. Rats in each group were sacrificed at Day 14 after the MCAO. (B) Western-blot analysis of NF-200 (n = 3), * p < 0.01, as compared with the AAV-empty group, # p < 0.05, as compared with the AAV-empty group. (C) Western-blot analysis of GAP-43 (n = 3), * p < 0.05, as compared with the AAV-empty group, # p < 0.01, as compared with the AAV-NT-1 group. (D) Western-blot analysis of MAP-2 (n = 3), * p < 0.05, as compared with the AAV-empty group. Index in PubMed under a CC BY license. PMID: 29487502

1. PubMed ID: 10.3109/09273948.2013.806989, Improved Retinal Ganglion Cell Survival through Retinal Microglia Suppression by a Chinese Herb Extract, Triptolide, in the DBA/2J Mouse Model of Glaucoma
2. PubMed ID: PMID:31966489, Gene silencing NMII promotes axonal regeneration against contusive spinal cord injury in rats
3. PubMed ID: 10.12659/MSM.900893, Changes in the Expression of miR-34a and its Target Genes Following Spinal Cord Injury In Rats

Visit bosterbio.com/anti-nefh-picoband-trade-antibody-a05307-boster.html to see all 17 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Nefh Antibody

For Research Use Only. Not for use in diagnostic procedures.