

Anti-Syntaxin 4/STX4 Antibody Picoband®

Catalog Number: A05345-1

About STX4

Enables sphingomyelin phosphodiesterase activator activity. Involved in several processes, including cornified envelope assembly; positive regulation of immune effector process; and positive regulation of protein localization. Located in several cellular components, including basolateral plasma membrane; cytoplasmic vesicle; and lamellipodium. Part of SNARE complex. Is active in glutamatergic synapse and postsynapse. Implicated in autosomal recessive nonsyndromic deafness.

Overview


Product Name	Anti-Syntaxin 4/STX4 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Syntaxin 4/STX4 Antibody Picoband® catalog # A05345-1. Tested in WB, IHC, ICC, IF, IP, Flow Cytometry, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q12846

Technical Details


Immunogen	E.coli-derived human Syntaxin 4/STX4 recombinant protein (Position: M1-I278). Human Syntaxin 4/STX4 shares 91.7% and 91.4% amino acid (aa) sequence identity with mouse and rat Syntaxin 4/STX4, respectively.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Human, Mouse, Rat Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human

	Immunoprecipitation, 0.5-2 ug/ml, Human Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5 ug/ml
--	--


Anti-Syntaxin 4/STX4 Antibody Picoband® (A05345-1) Images



Western blot analysis of Syntaxin 4/STX4 using anti-Syntaxin 4/STX4 antibody (A05345-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat liver tissue lysates, Lane 2: rat RH35 whole cell lysates, Lane 3: rat PC-12 whole cell lysates, Lane 4: mouse liver tissue lysates, Lane 5: mouse NIH/3T3 whole cell lysates, Lane 6: mouse HEPA1-6 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Syntaxin 4/STX4 antigen affinity purified polyclonal antibody (A05345-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for Syntaxin 4/STX4 at approximately 37 kDa. The expected band size for Syntaxin 4/STX4 is at 34 kDa.




IHC analysis of Syntaxin 4/STX4 using anti-Syntaxin 4/STX4 antibody (A05345-1). Syntaxin 4/STX4 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Syntaxin 4/STX4 Antibody (A05345-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of Syntaxin 4/STX4 using anti-Syntaxin 4/STX4 antibody (A05345-1). Syntaxin 4/STX4 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Syntaxin 4/STX4 Antibody (A05345-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


IF analysis of Syntaxin 4/STX4 using anti-Syntaxin 4/STX4 antibody (A05345-1). Syntaxin 4/STX4 was detected in an



immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-Syntaxin 4/STX4 Antibody (A05345-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Immunoprecipitating Syntaxin 4/STX4 in jurkat whole cell lysate. Western blot analysis of Syntaxin 4/STX4 using anti-Syntaxin 4/STX4 antibody (A05345-1). Lane 1: jurkat whole cell lysates (30ug), Lane 2: Rabbit control IgG instead of anti-Syntaxin 4/STX4 antibody in jurkat whole cell lysate, Lane 3: anti-Syntaxin 4/STX4 antibody (2ug) + jurkat whole cell lysate (500ug). After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-Syntaxin 4/STX4 antigen affinity purified polyclonal antibody (A05345-1) at a dilution of 0.5 ug/mL and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Syntaxin 4/STX4 at approximately 37 kDa. The expected band size for Syntaxin 4/STX4 is at 34 kDa.



Flow Cytometry analysis of K562 cells using anti-Syntaxin 4/STX4 antibody (A05345-1). Overlay histogram showing K562 cells stained with A05345-1 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-Syntaxin 4/STX4 Antibody (A05345-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Syntaxin 4/STX4 Antibody

For Research Use Only. Not for use in diagnostic procedures.