

888-466-3604 | support@bosterbio.com | www.bosterbio.com

### Anti-TLK1 Antibody

Catalog Number: A05635-1

### About TLK1

Rapidly and transiently inhibited by phosphorylation following the generation of DNA double-stranded breaks during S-phase. This is cell cycle checkpoint and ATM-pathway dependent and appears to regulate processes involved in chromatin assembly. Isoform 3 phosphorylates and enhances the stability of the t-SNARE SNAP23, augmenting its assembly with syntaxin. Isoform 3 protects the cells from the ionizing radiation by facilitating the repair of DSBs. In vitro, phosphorylates histone H3 at 'Ser-10'.

Yamakawa A., Gene 202:193-201(1997). Sillje H.H.W., EMBO J. 18:5691-5702(1999). Cabaniols J.-P., Mol. Biol. Cell 10:4033-4041(1999).

### Overview

Product Name	Anti-TLK1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-TLK1 Antibody (Catalog # A05635-1). Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UKI8

### **Technical Details**

Immunogen	Synthesized peptide derived from internal of human TLK1.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	lgG
Form	Liquid
Concentration	1 mg/ml



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com


Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope- specific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blotting: 1:500~1:3000



## BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

### Anti-TLK1 Antibody (A05635-1) Images



Western blot analysis of extracts from rat brain cells and NIH/3T3 cells, using TLK1 antibody A05635-1.

#### Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-TLK1 Antibody