

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-NARG1/NAA15 Antibody Picoband™

Catalog Number: A05636-1

About NAA15

NMDA receptor-regulated protein 1 (NARG1), also known as GA19 or Tbdn100 is a protein that in humans is encoded by the NAA15 gene. It is mapped to chromosome 4. NARG1 is the auxiliary subunit of the NatA (Nalpha-acetyltransferase A) complex. Both, Naa15 and Naa16 interact with the ribosome in yeast (via the ribosomal proteins, uL23 and uL29), humans and rat, thereby linking the NatA/Naa10 to the ribosome and facilitating co-translational acetylation of nascent polypeptide chains as they emerges from the exit tunnel. Furthermore, Naa15 might act as a scaffold for other factors, including the chaperone like protein HYPK (Huntingtin Interacting Protein K) and Naa50, the catalytic acetyltransferase subunit of NatE.

Overview

Product Name	Anti-NARG1/NAA15 Antibody Picoband™
Reactive Species	Human, Mouse
Description	Boster Bio Anti-NARG1/NAA15 Antibody Picoband [™] catalog # A05636-1. Tested in IHC, WB applications. This antibody reacts with Human, Mouse.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na $_2$ HPO $_4$, 0.05mg NaN $_3$.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9BXJ9

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human NARG1, different from the related mouse sequence by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Mouse, Human, By Heat Western blot, 0.1-0.5ug/ml, Human



888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-NARG1/NAA15 Antibody Picoband[™] (A05636-1) Images

130KD -	Figure 1. Western blot analysis of NARG1 using anti-NARG1
100KD	antibody (A05636-1).
70KD —	Electrophoresis was performed on a 5-20% SDS-PAGE gel at
55KD -	70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample
35KD-	under reducing conditions.
25KD-	lane 1: 293T whole cell lysates.
20112	After Electrophoresis, proteins were transferred to a
15KD -	Nitrocellulose membrane at 150mA for 50-90 minutes.
	Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5
	hour at RT. The membrane was incubated with rabbit anti-
	NARG1 antigen affinity purified polyclonal antibody (Catalog
	# A05636-1) at 0.5 ug/mL overnight at 4°C, then washed
	with TBS-0.1%Tween 3 times with 5 minutes each and
	probed with a goat anti-rabbit IgG-HRP secondary antibody
	at a dilution of 1:10000 for 1.5 hour at RT. The signal is
	developed using an Enhanced Chemiluminescent detection
	(ECL) kit (Catalog # EK1002) with Tanon 5200 system. A
	specific band was detected for NARG1 at approximately



Figure 2. IHC analysis of NARG1 using anti-NARG1 antibody (A05636-1).

101KD. The expected band size for NARG1 is at 101KD.

NARG1 was detected in paraffin-embedded section of mouse intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-NARG1 Antibody (A05636-1) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-NARG1/NAA15 Antibody ™