

Anti-NCOA7 Antibody (N-term)

Catalog Number: A05814-2

About NCOA7

NCOA7 enhances the transcriptional activities of several nuclear receptors. It is involved in the coactivation of different nuclear receptors, such as ESR1, THRB, PPARG and RARA.


Overview

Product Name	Anti-NCOA7 Antibody (N-term)
Reactive Species	Human, Mouse
Description	Boster Bio Anti-NCOA7 Antibody (N-term) (Catalog # A05814-2). Tested in WB, Flow Cytometry, IHC-P application(s). This antibody reacts with Human, Mouse.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8NI08


Technical Details

Immunogen	This NCOA7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 76-103 amino acids from the N-terminal region of human NCOA7.
Predicted Reactive Species	Bovine, Rat
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:1000 IHC-P: 1:10-1:50 FC: 1:10-1:50


Anti-NCOA7 Antibody (N-term) (A05814-2) Images




All lanes : Anti-NCOA7 Antibody (N-term) at 1:1000 dilution
Lane 1: HeLa whole cell lysate
Lane 2: human brain lysate
Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 106 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



NCOA7 Antibody (N-term) western blot analysis in HeLa, MDA-MB-453 cell line and mouse kidney tissue lysates (35ug/lane). This demonstrates the NCOA7 antibody detected the NCOA7 protein (arrow).



Formalin-fixed and paraffin-embedded human brain tissue reacted with NCOA7 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



NCOA7 Antibody (N-term) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NCOA7 Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.