

Anti-IDO-2/IDO2 Antibody Picoband®

Catalog Number: A06002-2

About IDO2

IDO2 (Indoleamine 2,3-dioxygenase 2), also called INDOLEAMINE 2,3-DIOXYGENASE-LIKE 1 or INDOL1, is an enzyme encoded by the INDOL1 gene which metabolizes tryptophan in the kynurenine pathway. By genomic sequence analysis, the INDOL1 gene is mapped on chromosome 8p12 just downstream of the INDO gene. And its exact cytogenetic location is 8p11.21. By database analysis using INDO as probe, followed by RT-PCR of total RNA from various tissues, IDO2 is cloned by human and mouse INDOL1. INDOL1 catabolizes tryptophan as determined by Kyn production, but unlike INDO, is inhibited by D-1-methyl-tryptophan (D-1MT) but not the L-1MT stereoisomer. The Gene Structure of the INDOL1 has 11 exons and spans 74 kb.

Overview


Product Name	Anti-IDO-2/IDO2 Antibody Picoband®
Reactive Species	Human, Monkey, Mouse, Rat
Description	Boster Bio Anti-IDO-2/IDO2 Antibody Picoband® catalog # A06002-2. Tested in ELISA, Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Monkey, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q6ZQW0

Technical Details


Immunogen	E.coli-derived human IDO-2/IDO2 recombinant protein (Position: M1-R357).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Mouse, Rat, Monkey Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5ug/ml, -


Anti-IDO-2/IDO2 Antibody Picoband® (A06002-2) Images



Western blot analysis of IDO-2/IDO2 using anti-IDO-2/IDO2 antibody (A06002-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions. Lane 1: rat liver tissue lysates, Lane 2: mouse liver tissue lysates, Lane 3: monkey kidney tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IDO-2/IDO2 antigen affinity purified polyclonal antibody (Catalog # A06002-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IDO-2/IDO2 at approximately 45KD. The expected band size for IDO-2/IDO2 is at 45KD.



Flow Cytometry analysis of K562 cells using anti-IDO-2/IDO2 antibody (A06002-2). Overlay histogram showing K562 cells stained with A06002-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-IDO-2/IDO2 Antibody (A06002-2, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.



IHC analysis of IDO-2/IDO2 using anti-IDO-2/IDO2 antibody (A06002-2). IDO-2/IDO2 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IDO-2/IDO2 Antibody (A06002-2) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IDO-2/IDO2 Antibody

For Research Use Only. Not for use in diagnostic procedures.