

Anti-Dio-1 DIDO1 Antibody

Catalog Number: A06019

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview

Product Name	Anti-Dio-1 DIDO1 Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Dio-1 DIDO1 Antibody catalog # A06019. Tested in ELISA, IHC, WB applications. This antibody reacts with Human, Mouse.
Application	ELISA, IF, IHC, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% stabilizing protein and 0.02% sodium azide. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9BTC0


Technical Details

Immunogen	The antiserum was produced against synthesized peptide derived from human DIDO1. AA range:161-210
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Suggested Dilutions

WB 1:500-1:2000
IHC 1:100-1:300
ELISA 1:10000
IF 1:50-200


Anti-Dio-1 DIDO1 Antibody (A06019) Images




Western Blot analysis of various cells using Dio-1 Polyclonal Antibody diluted at 1:2000



Western blot analysis of lysates from 293 cells, using DIDO1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using DNL1 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Anti-Dio-1 DIDO1 Antibody

For Research Use Only. Not for use in diagnostic procedures.